


Impact of inactivity on muscles more severe for older people

January 5 2018



Skeletal muscle tissue. Credit: University of Michigan Medical School

According to a recent study published in *The Journal of Physiology*, researchers have been able to document for the first time how the same period of inactivity has a greater and more severe impact on the muscle power of the lower limbs of the elderly than young people, which is essential for movements like climbing the stairs.

The disuse of muscles due to a sedentary lifestyle or short periods of inactivity caused by hospitalisation can dramatically enhance the decline in [muscle mass](#), [metabolic health](#) and [functional capacity](#). This loss of [muscle power](#) caused by disuse can be especially detrimental in the

elderly.

The research, conducted by the University of Udine in conjunction with the University of Padova, involved studying the impact of complete inactivity in a group of elderly subjects that were bedridden in a hospital environment for 2 weeks, and their results were compared with young subjects.

In the elderly subjects there was a difference in single muscle fibre response to disuse, a more pronounced loss of muscle mass and a change in how muscle contraction is controlled by the nervous system compared to young individuals. Furthermore the recovery phase was more difficult in the elderly group.

Carlo Reggiani, the lead investigator on the project commented on the findings: "While clinical and epidemiological data on inactivity in the elderly are abundant, experiments on disuse and inactivity are seldom performed in elderly for several reasons. The results obtained are relevant not only to understand the [inactivity](#)-dependent enhancement of the decline (in muscle mass, metabolic health and functional capacity) but also to design new rehabilitation protocols where timing and intensity of the sessions are optimized."

It is important to note that the study was carried out on healthy [elderly subjects](#) however the loss may be greater and the recovery more difficult in patients with diseases.

The researchers plan to follow-up the study by investigating the impairment of neural motor control on skeletal muscles, as an altered motor control was suggested when measuring the electrical activity of the muscles.

More information: Enrico Rejc et al, Loss of maximal explosive

power of lower limbs after two weeks of disuse and incomplete recovery after retraining in older adults, *The Journal of Physiology* (2017). [DOI: 10.1113/JP274772](https://doi.org/10.1113/JP274772)

Provided by The Physiological Society

Citation: Impact of inactivity on muscles more severe for older people (2018, January 5)
retrieved 23 April 2024 from
<https://medicalxpress.com/news/2018-01-impact-inactivity-muscles-severe-older.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.