

Study examines differences in hip fracture rates among nursing homes

22 January 2018

In a nationally representative study, researchers found considerable variation in the rates of hip fractures across US nursing home facilities. The *Journal of the American Geriatrics Society* study identified a number of modifiable facility-level characteristics that might be addressed, but the majority of the variation in the risk of hip fracture remained unexplained.

Although the Centers for Medicare and Medicaid Services have determined that the proportion of residents that experience injurious falls is an important nursing home quality measure, the authors caution against the use of hip fracture rates as a similar quality measure.

"Although much of the variation was unexplained by measured characteristics in our study, psychotropic medication prescribing and staffing requirements emerged as plausible targets for initiatives to reduce [hip fracture](#) rates in nursing home facilities," said lead author Dr. Andrew Zullo, of Brown University School of Public Health.

More information: Andrew R. Zullo et al, Facility and State Variation in Hip Fracture in U.S. Nursing Home Residents, *Journal of the American Geriatrics Society* (2018). [DOI: 10.1111/jgs.15264](https://doi.org/10.1111/jgs.15264)

Provided by Wiley

APA citation: Study examines differences in hip fracture rates among nursing homes (2018, January 22) retrieved 24 October 2021 from <https://medicalxpress.com/news/2018-01-differences-hip-fracture-nursing-homes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.