

Yazidi women suffering from high percentage of C-PTSD, new study finds

January 31 2018

As the Islamic State of Iraq and Syria (ISIS) stormed Iraq and attempted to conquer the country in 2014, its members committed genocide against the Yazidi population, a Kurdish religious minority. Men were systematically executed. Women were captured, forced into sexual slavery and repeatedly raped, beaten, sold and locked away.

What are the long-term psychological effects of the conditions these women endured? Following severe trauma, people may develop post traumatic stress disorder (PTSD). Recently, specific diagnostic criteria were suggested for another psychiatric disorder following trauma, i.e., complex post-traumatic stress disorder (C-PTSD). Whereas PTSD typically occurs following a single traumatic event, C-PTSD is typically associated with prolonged trauma where one's destiny is under another's control and escape - from captivity, for example—is unfeasible. As opposed to PTSD, which can be triggered by trauma reminders, C-PTSD is conceived to be a more deeply-rooted disorder that affects the very core of one's self-organization.

A comprehensive study led by Bar-Ilan University researchers, just published in *World Psychiatry*, has shown that a very high percentage of female Yazidi former ISIS captives were suffering from C-PTSD (>50%) in addition to others with PTSD (23%). The women, all in their twenties, were surveyed over a two-month period in four refugee camps. "This main finding of over 50% C-PTSD prevalence cannot be emphasized enough, as PTSD and C-PTSD require different therapeutic interventions," says Dr. Yaakov Hoffman, of the Interdisciplinary

Department of Social Sciences at Bar-Ilan University, who led the study. Focusing solely on PTSD interventions is inadequate. With that in mind, the Bar-Ilan University team is planning to set up a training program for Kurdish mental health workers.

Another important finding of the research is the greater sensitivity of victims with C-PTSD to post ISIS conditions. Although ISIS no longer poses a threat to them, they still feel less safe, and their current living conditions may exacerbate their symptoms. Hoffman therefore emphasizes that "NGO's working with such victims need to be particularly sensitive not only to the unique Yazidi culture but also to the creation of an environment that is perceived by the former captives as protective, both on objective and subjective levels".

This study required extensive cooperation from three different countries (Israel, Germany and Kurdistan) by a multidisciplinary group from psychology, psychiatry, life sciences, brain science, Arabic, and communications. In addition to the Bar-Ilan group leading this study, the team included authors from Ariel and Haifa Universities, as well as researchers from Luftbrücke Irak, Germany, and the Erbil Psychiatric Hospital/Emma Organization for Human Development in Kurdistan. The authors are in the midst of continued Israeli-Kurd academic cooperation, both in the formation of suitable training programs as well as additional research designed to illuminate basic issues relevant to interventions on the ground.

More information: Yaakov S.G. Hoffman et al, Complex PTSD and its correlates amongst female Yazidi victims of sexual slavery living in post-ISIS camps, *World Psychiatry* (2018). [DOI: 10.1002/wps.20475](https://doi.org/10.1002/wps.20475)

Provided by Bar-Ilan University

Citation: Yazidi women suffering from high percentage of C-PTSD, new study finds (2018, January 31) retrieved 18 April 2024 from <https://medicalxpress.com/news/2018-01-yazidi-women-high-percentage-c-ptsd.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.