

Ibuprofen can damage men's endocrine systems

13 February 2018



Ibuprofen could reduce testosterone production in men.
Credit: Wikipedia

Ibuprofen can reduce the production of testosterone in men, shows new research.

Taking the painkiller daily for just two weeks resulted in a [hormone](#) imbalance among young male participants that more closely resembled that of older men whose testes produce less [testosterone](#), say the scientists behind the study.

"It's worrying because it's important that men's testosterone levels are stable. Otherwise, it can lead to problems with, for example, muscle development, mood, and potency," says David Møbjerg Kristensen, a senior researcher at the Danish Headache Center at Rigshospitalet and co-author of the new study, which is published in the scientific journal *PNAS*.

Ibuprofen knocks endocrine system out of balance

Kristensen and colleagues conducted both a randomised control trial and laboratory experiments, which showed that [ibuprofen](#) can reduce the testes' ability to create testosterone.

In the randomised trial, 31 men between the age of 18 and 35 were split into two groups. One was given 600 milligrams of ibuprofen every day for 44 days. The other group received a placebo.

Throughout the trial, scientists measured the content of luteinizing hormone (lutropin) in the participants' blood samples.

Lutropin indicates how much testosterone is produced by the testes, says Kristensen.

When less testosterone is produced, the brain tries to compensate by emitting large amounts of lutropin. This sends a message to the testes to produce more testosterone. So, elevated levels of lutropin in the blood indicates lower testosterone production.

The 14 male trial participants who received ibuprofen all had elevated levels of lutropin after just two weeks.

Endocrine system resembled that of a 70-year-old

When the brain tries to compensate for reduced testosterone levels by emitting lutropin, it is called compensatory hypogonadism.

"Compensatory hypogonadism is normally observed among older men, because as the years go by the testes being to produce less testosterone," says Kristensen.

"In our experiment we see that young men who take ibuprofen resemble what we see in 70-year-old men. We don't yet know what the consequences could be, and whether the conditions persists after you stop taking ibuprofen. Subsequent studies will look into this," he says.

Testes produce less testosterone

As part of the study, Kristensen's colleagues analysed testicular tissue to study how the testes responded to ibuprofen. To do this, they studied tissue from coma patients who had a complete loss of brain function.

These laboratory experiments showed that the testes precipitation of testosterone was reduced when men received ibuprofen.

But it was not just testosterone: All hormones produced in the testes were affected by ibuprofen.

First study of its kind

Professor Cecilia Ramlau-Hansen from the Department of Public Health at Aarhus University, Denmark, describes the study as "exciting" and "well executed." She was not involved in the research.

"The scientists use a range of approaches to investigate how ibuprofen influences the balance of hormones and the results of their research support each other," says Ramlau-Hansen.

One of the strengths of the study is the randomised trial, in which the male participants are split into two groups: One receiving ibuprofen, the other receiving a placebo. Randomised trials normally lead to good results, she says.

"As far as I know it's the first time anyone has made a randomised control trial to test how ibuprofen effects hormone balance," says Ramlau-Hansen.

Over-the-counter medicine can also have side effects

However, one drawback is that the study is not especially large with just 31 participants.

Moreover, the new study does not investigate how ibuprofen influences the hormone balance in the long term. They do not know whether the hormone imbalance returns to normal after taking ibuprofen for a certain period, or whether the hypogonadism becomes permanent.

The results indicate that men should use ibuprofen

sparingly, even if it is available to purchase over-the-counter and without prescription, say Ramlau-Hansen and Kristensen.

Some people need to take ibuprofen to treat pain, such as arthritis patients. But if you have a sports injury or headache, you should consider not taking ibuprofen.

"Research shows that many athletes, both professional and amateurs, take ibuprofen if they have an injury but wish to continue playing anyway," says Kristensen.

But the new study indicates that it might be better to take a break until the injury is healed. When you exercise, it is especially important to have a balance of testosterone, which plays an important role in [muscle development](#), he says.

Previous studies reveal mixed results

Kristensen and colleagues previously showed that boys have an increased risk of birth defects of their sexual organs if their mother took paracetamol painkillers while pregnant.

Other research into painkillers and [hormone imbalance](#) produced mixed results, says Ramlau-Hansen.

Some studies have shown that painkillers, such as ibuprofen and paracetamol, can disturb the hormone balance and reduce the testes production of testosterone. Other studies have indicated that painkillers can reduce sperm quality.

Another large population study by Ramlau-Hansen did not find any connection between painkillers and a reduced [testosterone level](#).

More information: David Møbjerg Kristensen et al. Ibuprofen alters human testicular physiology to produce a state of compensated hypogonadism, *Proceedings of the National Academy of Sciences* (2018). [DOI: 10.1073/pnas.1715035115](https://doi.org/10.1073/pnas.1715035115)

This story is republished courtesy of [ScienceNordic](#), the trusted source for English-language science

news from the Nordic countries. Read the original story [here](#).

Provided by ScienceNordic

APA citation: Ibuprofen can damage men's endocrine systems (2018, February 13) retrieved 20 August 2018 from <https://medicalxpress.com/news/2018-02-ibuprofen-men-endocrine.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.