

Higher biopsy rates for women undergoing screening breast MRI

February 14 2018



(HealthDay)—Women undergoing screening breast magnetic resonance

imaging (MRI) have higher biopsy rates and significantly lower cancer yield findings compared with screening mammography alone, regardless of personal history of breast cancer (PHBC), according to a study published online Feb. 12 in *JAMA Internal Medicine*.

Diana S.M. Buist, Ph.D., M.P.H., from the Kaiser Permanente Washington Health Research Institute in Seattle, and colleagues conducted an observational cohort study of six Breast Cancer Surveillance Consortium registries. Data were included for a sample of 812,164 women undergoing a total of 2,048,994 digital [mammography](#) and/or breast MRI screening episodes from 2003 through 2013. Outcomes were stratified by PHBC.

The researchers found that for women with PHBC, age-adjusted core and surgical biopsy rates were doubled after MRI versus mammography (57.1 versus 23.6 per 1,000 episodes). The differences were even larger for women without PHBC (84.7 and 14.9 per 1,000 episodes, respectively). In women with PHBC, the ductal carcinoma in situ and invasive biopsy yield was significantly higher after mammography versus MRI episodes (404.6 versus 267.6 per 1,000 episodes); for women without PHBC, the yield was nonsignificantly higher (279.3 versus 214.6 per 1,000 episodes). Regardless of PHBC, high-risk benign lesions were more commonly identified after MRI.

"Further work is needed to identify [women](#) who will benefit from screening MRI to ensure an acceptable benefit-to-harm ratio," the authors write.

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

Citation: Higher biopsy rates for women undergoing screening breast MRI (2018, February 14)
retrieved 19 September 2024 from

<https://medicalxpress.com/news/2018-02-higher-biopsy-women-screening-breast.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.