

Do pain medications carry different heart risks?

22 February 2018



have similar cardiovascular safety profiles," said co-author Dr. Chia-Hsuin Chang, of the National Taiwan University Hospital.

More information: Xiaogang Liu et al, Ticagrelor attenuates myocardial ischemia-reperfusion injury possibly through downregulating galectin-3 expression in the infarct area of rats, *British Journal of Clinical Pharmacology* (2018). DOI: [10.1111/bcp.13536](https://doi.org/10.1111/bcp.13536)

Provided by Wiley

Credit: CC0 Public Domain

Prior studies have suggested that non-steroidal anti-inflammatory drugs (NSAIDs) may be linked with higher cardiovascular risks, but few have assessed potential different cardiovascular risk between NSAID classes or across individual NSAIDs.

A *British Journal of Clinical Pharmacology* study including 55,629 [patients](#) with hypertension in a Taiwanese database did not observe different cardiovascular risks in patients who received cyclooxygenase-2 enzyme selective NSAIDs or nonselective NSAIDs during 4 weeks of follow-up.

The study also found no apparent difference in cardiovascular risk when comparing celecoxib with diclofenac, ibuprofen, or naproxen, although a significantly increased risk was observed when comparing celecoxib with mefenamic acid.

"Our results provide important information about the comparative safety of alternative NSAID use in patients with hypertension in real-world settings. Under low-to-moderate daily dose and a short-term treatment period, most commonly used NSAIDs

APA citation: Do pain medications carry different heart risks? (2018, February 22) retrieved 31 October 2020 from <https://medicalxpress.com/news/2018-02-pain-medications-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.