

Study sheds light on how childhood RSV can lead to asthma

March 2 2018, by Christina Echegaray

Infants who have higher amounts of the bacterium *Lactobacillus* present in their nose or upper part of the throat during an acute respiratory syncytial virus (RSV) infection are less likely to develop childhood wheezing later in life, a new Vanderbilt-led Center for Asthma Research study found.

Lactobacillus is a type of "friendly" bacteria that normally live in the digestive, urinary, and genital systems without causing disease.

The study was published in the *Journal of Allergy and Clinical Immunology*. The latest finding in the research on RSV, one of the most common causes of upper and lower [acute respiratory infection](#) in young children worldwide, is important for researchers as they look to create interventions that prevent the development of [childhood](#) wheezing illnesses, including asthma.

Childhood wheezing illnesses have commonly been associated with RSV, however, the pathways underlying the association aren't fully understood.

"These findings provide new insights not only about how RSV infection in infancy may cause wheezing later in life, but also how the normal bacteria that live in and on us may protect us from disease," said Christian Rosas-Salazar, MD, MPH, one of the study's first authors and assistant professor of pediatrics.

The researchers conducted a study of previously healthy, term infants

who had laboratory-confirmed RSV acute respiratory infection (ARI).

The infants were enrolled in the INSPIRE (Infant Susceptibility to Pulmonary Infections and Asthma Following RSV Exposure) study—a population-based birth cohort of infants born between June and December, designed so that the first RSV infection during infancy could be studied. Eligible infants were enrolled from collaborating general pediatric practices throughout the Middle Tennessee region.

Researchers collected samples of bacteria from the infants' noses and upper throats. The infants were followed until age 2, when their wheezing outcomes were assessed. The detection and increased abundance of *Lactobacillus* in the infants' noses and throats was consistently higher in infants who did not develop wheezing at 2 years.

With this information, the researchers suggested, "the detection of *Lactobacillus* in the nasopharynx of RSV-infected [infants](#) could be used as a biomarker for the later development of childhood wheezing illnesses."

Provided by Vanderbilt University

Citation: Study sheds light on how childhood RSV can lead to asthma (2018, March 2) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-03-childhood-rsv-asthma.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
