


Detecting diminished dopamine-firing cells inside brain could reveal earliest signs of Alzheimer's

March 27 2018



PET scan of a human brain with Alzheimer's disease. Credit: public domain

A new link between diminished input from dopamine-firing cells deep

inside the brain and the ability to form new memories could be crucial in detecting the earliest signs of Alzheimer's disease.

Scientists from the National Institute for Health Research (NIHR) Sheffield Biomedical Research Centre at the University of Sheffield have discovered a loss of cells that use dopamine - a neurotransmitter that has a number of functions including regulating movement and emotional responses - may cause the part of the brain responsible for forming [new memories](#) to function less effectively.

The findings could revolutionise screening for the early signs of Alzheimer's disease - which affects more than 520,000 people in the UK - changing the way brain scans are acquired and interpreted as well as using different memory tests.

Lead author of the study, Professor Annalena Venneri, from the Sheffield Institute for Translational Neuroscience (SITraN) at the University of Sheffield, said: "Our findings suggest that if a small area of brain cells, called the [ventral tegmental area](#), does not produce the right amount of dopamine for the hippocampus, a small organ located within the brain's temporal lobe, it will not work efficiently.

"The hippocampus is associated with forming new memories, therefore these findings are crucial to the early detection of Alzheimer's disease. The results point at a change which happens very early on, which might trigger Alzheimer's disease.

"This is the first study to demonstrate such a link in humans."

Professor Venneri and fellow lead author Dr Matteo De Marco acquired 3Tesla Magnetic Resonance Imaging (MRI) scans on 51 healthy adults, 30 patients with a diagnosis of mild cognitive impairment, and 29 patients with a diagnosis of Alzheimer's disease. 3Tesla MRIs are twice

the normal strength of normal MRI scans generating the highest quality images.

The results showed a key link between the size and function of the ventral tegmental area, the size of the hippocampus and the ability to learn new material.

"More studies are necessary, but these findings could potentially lead to a new way of screening the elderly population for early signs of Alzheimer's disease, changing the way brain scans are acquired and interpreted and using different memory tests," said Professor Venneri who is also an Honorary Consultant at Sheffield Teaching Hospital NHS Foundation Trust.

"Another possible benefit is that it might lead to a different treatment option with the potential to change or halt the course of the disease very early, before major symptoms manifest.

"We now want to establish how early alterations in the ventral tegmental area can be seen and also test whether these alterations can be counteracted with treatments already available."

The results of the study are published today (27 March 2018) in the *Journal of Alzheimer's Disease*.

Provided by University of Sheffield

Citation: Detecting diminished dopamine-firing cells inside brain could reveal earliest signs of Alzheimer's (2018, March 27) retrieved 19 September 2024 from <https://medicalxpress.com/news/2018-03-diminished-dopamine-firing-cells-brain-reveal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.