

Unraveling the immunopathogenesis of Johne's disease

April 2 2018




One of the cattle with Johne's disease showing significant weight loss. Photo provided by National Agriculture and Food Research Organization. Credit: Hokkaido University

Researchers of Hokkaido University, the National Agriculture and Food Research Organization (NARO), and Tohoku University demonstrated that a physiologically active substance called prostaglandin E2 (PGE2) impairs the immune response by upregulating the expression of an immunoinhibitory molecule, programmed death-ligand 1 (PD-L1), in cattle affected with Johne's disease. The discovery is expected to help

develop a method to control Johne's disease, which is most frequently reported among legally designated infectious bovine diseases in Japan.

Johne's [disease](#) is a type of chronic enteritis caused by *Mycobacterium avium* subsp. *paratuberculosis*, leading to diarrhea and weight loss in [cattle](#), and occasionally death. This disease has led to endemics in Japan and 816 clinical cases were reported in 2017. Cattle diagnosed with the disease must be culled to prevent its spread. Cattle farmers are also required to give regular check-ups to all other cattle at the same farm for a certain period, disinfect their cattle barns and are not allowed to move animals out of the farm freely. There has been no effective method to prevent the spread of the disease, including vaccination.

The researchers investigated the immunosuppressive effects of PGE₂, which is known to appear in increased amounts in cattle with Johne's disease. Their experiments showed PGE₂ has immunosuppressive effects on T-cell, a type of lymphocyte that plays an important role in the immune system. PGE₂ also induced the upregulation of PD-L1, a immunosuppressive molecule, in [immune cells](#) isolated from cows. The researchers also found PGE₂ and PD-L1 were co-expressed in intestinal lesions of the infected cattle.



Immunohistochemical analysis showing expression of PGE2 (left) and PD-L1 (right) in the ileal mucosa of a cow with Johne's disease. Credit: Hokkaido University

In addition, they found that the inhibition of cyclooxygenase-2 (COX-2), which is involved in the synthesis of PGE2, boosted immune response to the mycobacteria causing Johne's disease. When COX-2 inhibitor is combined with anti-PD-L1 blocking antibodies, it further enhances the [immune response](#) to the mycobacteria.

"Our data suggests that immunosuppressive effects of PGE2 are strongly connected to the progression of Johne's disease," says Associate Professor Satoru Konnai of the research team at Hokkaido University. "We plan to conduct clinical trials to verify how the COX-2 inhibitor and anti-PD-L1 antibodies boost immune responses in cattle with Johne's disease, in addition to examining the roles of PGE2 in cattle afflicted with other diseases."

More information: Yamato Sajiki et al. Prostaglandin E2 induction suppresses the Th1 immune responses in cattle with Johne's disease, *Infection and Immunity* (2018). [DOI: 10.1128/IAI.00910-17](https://doi.org/10.1128/IAI.00910-17)

Provided by Hokkaido University

Citation: Unraveling the immunopathogenesis of Johne's disease (2018, April 2) retrieved 20 April 2024 from <https://medicalxpress.com/news/2018-04-unraveling-immunopathogenesis-johne-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.