

Pre-op chronic opioid use ups poor outcomes post spinal fusion

2 April 2018



department visits, readmission, and continued postoperative use. Additionally, one year after surgery, long-term opioid users had an increased risk for epidural/facet joint injections, revision fusion, and new onset constipation.

"Preoperative chronic opioid therapy is a modifiable risk factor for complications, [readmission](#), [adverse events](#), and increased costs after one- and two-level posterior lumbar [fusion](#)," the authors write.

More information: [Abstract/Full Text](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

(HealthDay)—Preoperative chronic opioid use is associated with poor outcomes and continued dependence after posterior lumbar fusion, according to a study published online March 20 in *Spine*.

Nikhil Jain, M.D., from The Ohio State University in Columbus, and colleagues used commercial insurance data (from 2007 to Q3-2015) to evaluate preoperative [opioid](#) use in 24,610 patients undergoing primary one- and two-level posterior lumbar fusion. The authors also assessed associated 90-day complications, emergency department visits, readmissions, one-year adverse events, and costs.

The researchers found that 22.3 percent of patients had documented opioid use for more than six months before surgery, and 87.4 percent of these had continued long-term use postoperatively. Preoperative chronic opioid use was associated with higher risk for 90-day wound complications, pain diagnoses, emergency

APA citation: Pre-op chronic opioid use ups poor outcomes post spinal fusion (2018, April 2) retrieved 26 September 2021 from <https://medicalxpress.com/news/2018-04-pre-op-chronic-opioid-ups-poor.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.