

Patients with high-risk clinical features are at high risk for acute aortic dissection

April 10 2018

Aortic dissection: high risk clinical features
Case-control study: patients >18y, acute atraumatic truncal pain
Cases n=194, controls n=776

	Tearing/ripping pain Sens 13.6%, Spec 99.7% LR- 0.87, LR+ 42.07	New murmur Sens 20.6%, Spec 97.8% LR- 0.81, LR+ 9.41	
	Pulse deficit Sens 20.6%, Spec 99.3% LR- 0.8, LR+ 31.14	Chest Xray* Sens 52.4%, Spec 94.9% LR- 0.5, LR+ 10.23 Widened mediastinum /absent aortic notch	
	Hypotension Sens 22.2%, Spec 98.7% LR- 0.79, LR+ 17.2	Abrupt-Onset pain Sens 95.9%, Spec 61.3% LR- 0.07, LR+ 2.48	

Ohle 2018 doi 10.1111/acem.13356



Aortic dissection: high-risk clinical features. Credit: Kirsty Challen, B.Sc., MBChB, MRes, Ph.D., Lancashire Teaching Hospitals, United Kingdom.

Patients with one or more high-risk clinical features (tearing pain, hypotension, pulse deficit, neurologic deficit, new murmur) should be considered high risk for acute aortic dissection (AAD). That is the finding of a study to be published in the April 2018 issue of *Academic Emergency Medicine (AEM)*, a journal of the Society for Academic Emergency Medicine (SAEM).

The lead author of the study is Dr. Robert Ohle, MSc, MA, MB, FRCPC, Department of Emergency Medicine, the Ottawa Hospital Research Institute, University of Ottawa, Ottawa, Ontario.

The study by Ohle, et al, further found that patients with no high-risk and multiple low-risk features (absence of abrupt-onset pain, history of ischemic heart disease and diabetes) are at low risk for acute aortic dissection.

A novel finding of the study was that a history of [ischemic heart disease](#) decreased the probability of AAD.

The study suggested that educational strategies focusing on disseminating the diagnostic accuracy of clinical findings could reduce time to diagnosis and that further research should focus on the ability of a combination of these factors in the assessment of a patient for AAD.

The findings support the need for prospective data collection to address the reproducibility of subjective historical features and the accuracy of D-dimer in a lower-risk population.

Lane McNeil Smith, MD, PhD, assistant professor, Lexington Emergency Medicine, Wake Forest Baptist Health, commented:

"Ohle et al have done a great job at telling us how [acute aortic dissection](#) presents to [emergency](#) departments. Are we finally on the verge of a

decision rule for this disease that improves our miss rate without ballooning non-therapeutic imaging and costs? This study is a step in the right direction and the authors are in a good position to find the right balance of improved accuracy and increased cost."

More information: Robert Ohle et al, High Risk Clinical Features for Acute Aortic Dissection: A Case-Control Study, *Academic Emergency Medicine* (2017). [DOI: 10.1111/acem.13356](https://doi.org/10.1111/acem.13356)

Provided by Society for Academic Emergency Medicine

Citation: Patients with high-risk clinical features are at high risk for acute aortic dissection (2018, April 10) retrieved 20 September 2024 from <https://medicalxpress.com/news/2018-04-patients-high-risk-clinical-features-high.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.