

Blood test may predict future risk of cardiovascular events

10 April 2018

Despite heart disease and type 2 diabetes being among the leading causes of death in the U.S., the mechanisms leading to and linking these two diseases remain incompletely understood. A new study by investigators at Brigham and Women's Hospital may help shed light on a molecular pathway that heart disease and diabetes share, and points to a biomarker that is elevated in women at risk of cardiovascular disease and diabetes. The team's results were presented by Deirdre Tobias, ScD, at the AHA Epidemiology and Lifestyle meeting and simultaneously published in *Circulation: Genomic and Precision Medicine*.

"We examined more than 27,000 women in the Women's Health Study and found that a one-time measurement of branched chain amino acids in the blood stream - a test that now can be easily done - predicted future risk of [cardiovascular events](#) to the same extent and independent of LDL cholesterol and other risk factors," said corresponding author Samia Mora, MD, of the Center for Lipid Metabolomics at BWH. "This was particularly so for women who developed type 2 diabetes prior to their cardiovascular disease."

Branched chain amino acids (BCAAs) are thought to play a causal role in the development of insulin resistance and type 2 diabetes. However, few studies have evaluated the association between BCAAs and [cardiovascular disease](#). To investigate, the team measured BCAA levels in blood samples using NMR spectrometry. Of the more than 27,000 women studied, 2,207 experienced a cardiovascular event over the 18 year follow up period.

The team found a positive association between BCAA levels and incident of CVD events. This association was much more pronounced in [women](#) who developed diabetes before experiencing a cardiovascular event. The team adjusted for other biomarkers related to diabetes - including hbA1c - finding evidence that BCAAs may be tied to

downstream biomarkers of type 2 [diabetes](#) metabolism.

"Impaired BCAA metabolism may represent a shared pathway of the metabolic pathophysiology that links the risks of T2D and CVD," the authors conclude.

"There is little known at this time as to what leads to elevated levels of BCAAs or what can be done clinically to reduce them, and if this leads to a reduction in risk, but further research will target these important questions," said Tobias.

More information: Tobias DK et al. "Circulating Branched-Chain Amino Acids and Incident Cardiovascular Disease in a Prospective Cohort of US Women" *Circulation: Genomic and Precision Medicine* DOI: [10.1161/CIRCGEN.118.002157](https://doi.org/10.1161/CIRCGEN.118.002157)

Provided by Brigham and Women's Hospital

APA citation: Blood test may predict future risk of cardiovascular events (2018, April 10) retrieved 12 May 2021 from <https://medicalxpress.com/news/2018-04-blood-future-cardiovascular-events.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.