

Starting aerobic exercise soon after concussion improves recovery time, study finds

20 April 2018



Exercising on stationary bikes at U of T's David L. MacIntosh Sport Medicine Clinic, located in the Goldring Centre for High Performance Sport. Credit: Faculty of Kinesiology & Physical Education

A new study from the University of Toronto's Faculty of Kinesiology & Physical Education has found that starting aerobic exercise sooner rather than later after a diagnosed concussion contributes to a faster recovery and return to sport, school and work.

The study, published in the journal *PLOS ONE*, supports the view that aerobic exercise is safe and potentially protective in symptomatic individuals.

"We knew aerobic exercise had a positive influence in other conditions that affect the brain (such as depression, stroke and cognitive impairment) and in cases of persistent symptoms following concussion. However, only a couple of studies previously examined aerobic exercise in the acute period, and none as early as we did," said Dr. David Lawrence, staff physician at the David L. MacIntosh Sport Medicine Clinic at U of T

and principal author of the study.

So, how soon after a concussion is it safe to start doing aerobic exercise? Lawrence says the study, which followed the recovery of 253 people between the ages of 15 and 20, has shown that some individuals benefit from starting low-impact, [aerobic activity](#) as early as 24 hours after injury.

For each successive day of delaying the start of aerobic exercise, individuals had a less favourable recovery trajectory, according to the study. Initiating aerobic exercise at three and seven days following injury was associated with a reduced probability of 36.5 per cent and 73.2 per cent respectively of a faster full return to sport, and a reduced probability of 45.9 per cent and 83.1 per cent respectively of a faster full return to school and work.

"Historically, [concussion management](#) was based on a simple recipe of rest until your symptoms go away. However, what we have realized is that in many people symptoms take time to resolve and prolonged periods of rest may have a negative impact because rest was interpreted as no activity in sport, school, work, screen and social activities," said Michael Hutchison, director of the concussion program at the MacIntosh clinic and co-author of the study.

"We still believe that a brief reduction in activity from normal levels is beneficial," he said.

"However, we recommend maintaining activity levels that do not exacerbate symptoms in the acute period."

Lawrence suggests initially doing low-impact aerobic exercise with minimal head movement, such as stationary cycling, elliptical and walking. Jogging and swimming are not recommended at this stage as they involve greater head movement.

"Exercise recommendations should be individualized based on the clinical assessment, but we feel the intensity of [aerobic exercise](#) should begin with low-intensity (you should be able to have a conversation with someone as you exercise) and should not worsen the symptoms," says Lawrence.

The results of this study did not observe a different recovery pattern between men and women. However, a history of concussion, higher symptom burden and loss of consciousness were associated with a prolonged recovery.

The scientists recommend any exercise plans prescribed by physicians following an injury such as a [concussion](#) should include recommendations about when to start exercise post-injury, in addition to the customary recommendations about the frequency, intensity and type of [exercise](#).

More information: David Wyndham Lawrence et al. Earlier time to aerobic exercise is associated with faster recovery following acute sport concussion, *PLOS ONE* (2018). [DOI: 10.1371/journal.pone.0196062](#)

Provided by University of Toronto

APA citation: Starting aerobic exercise soon after concussion improves recovery time, study finds (2018, April 20) retrieved 21 September 2019 from <https://medicalxpress.com/news/2018-04-aerobic-concussion-recovery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.