

Targeted radiotherapy for breast cancer offers good quality of life and fewer side effects

April 22 2018

Quality of life for women treated with a more targeted radiotherapy treatment - called accelerated partial breast irradiation - is at least as good as quality of life for women treated with standard radiotherapy, according to research presented at the ESTRO 37 conference and published simultaneously in *The Lancet Oncology*.

The [treatment](#) uses small tubes to deliver a higher dose of [radiotherapy](#) directly to the site of the tumour over a shorter period of time. Previous research shows that it is just as effective at treating early stage [breast cancer](#) as radiotherapy to the whole of the breast.

Researchers say the targeted treatment, which takes a week, is a good option for [women](#) with early breast cancer, especially those who would struggle with the three to six-week schedule of traditional radiotherapy.

The analysis of quality of life data was led by Dr Rebekka Schäfer, a clinician in the department of radiation oncology at University Hospital Würzburg, Germany. It involved more than 600 women treated at 16 hospitals across Europe who all had surgery to remove an early stage tumour.

Around half of the women received whole breast radiotherapy while the other half received the accelerated partial breast irradiation treatment. The research was part of a larger study that has already shown that the

treatments are equally good in terms of survival and preventing cancer recurring in the breast.

Dr Schäfer said: "This treatment focuses only on the area of the breast where the tumour has been removed and so it offers a number of advantages. It allows us to deliver a higher dose of radiation to a smaller area, which means the course of treatment is shortened from three to six weeks down to one week or less. It also makes it easier to avoid unnecessary radiation to other parts of the body, such as the heart and lungs.

"However, compared to external radiotherapy this is a more invasive treatment, so we wanted to see how it affected women's quality of life and their symptoms, both when they are treated and in the longer term."

Researchers questioned women about their symptoms and quality of life before and after their radiotherapy treatments, and at regular intervals for up to five years following treatment.

They found that women's quality of life was equally good in both groups and remained so over the five years.

Symptoms of breast swelling and pain, and skin problems of the breast were worse in women treated with whole breast radiotherapy, especially immediately after treatment and three months later. Swelling and discomfort in the arm was also somewhat worse in the women given standard treatment.

Dr Schäfer added: "More and more women are surviving breast cancer for longer and longer, so we need to be sure that treatments do not detract from their quality of life in the long term.

"Research like this is vital because it allows us to offer treatment

choices. For a variety of reasons patients may find it difficult to manage a treatment schedule that lasts up to six weeks. They may have to travel a long way to reach hospital, or they may wish to return to work as soon as possible, so they might prefer a treatment that can be carried out in a week or less."

The researchers plan a ten-year follow-up of women in the trial to see if [quality](#) of life remains good and to see if there is any difference in cancer recurrence between the two treatments.

President of ESTRO, Professor Yolande Lievens, head of the department of radiation oncology at Ghent University Hospital, Belgium, said: "Until recently, radiotherapy to the whole breast has been the gold-standard treatment following surgery for early stage breast cancer. However, different technical approaches to partial breast irradiation have been developed and are gaining acceptance in clinical practice. The treatment described is already available at many hospitals in Europe. This study adds to evidence that accelerated partial breast treatments may be a valuable alternative for women with early stage breast cancer."

More information: Rebekka Schäfer et al, Quality-of-life results for accelerated partial breast irradiation with interstitial brachytherapy versus whole-breast irradiation in early breast cancer after breast-conserving surgery (GEC-ESTRO): 5-year results of a randomised, phase 3 trial, *The Lancet Oncology* (2018). [DOI: 10.1016/S1470-2045\(18\)30195-5](https://doi.org/10.1016/S1470-2045(18)30195-5)

Provided by European Society for Radiotherapy and Oncology

Citation: Targeted radiotherapy for breast cancer offers good quality of life and fewer side effects (2018, April 22) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2018-04-radiotherapy-breast-cancer-good-quality.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.