


Noninvasive spinal stimulation method enables paralyzed people to regain use of hands, study finds

April 26 2018, by Stuart Wolpert



Credit: AI-generated image ([disclaimer](#))

The ability to perform simple daily tasks can make a big difference in people's lives, especially for those with spinal cord injuries. A UCLA-led team of scientists reports that six people with severe spinal cord injuries—three of them completely paralyzed—have regained use of

their hands and fingers for the first time in years after undergoing a nonsurgical, noninvasive spinal stimulation procedure the researchers developed.

At the beginning of the study, three of the participants could not move their fingers at all, and none could turn a doorknob with one hand or twist a cap off a plastic water bottle. Each of them also had great difficulty using a cellphone. After only eight researcher-led training sessions with the spinal stimulation, all six individuals showed substantial improvements. The study participants had chronic and severe paralysis for more than one year, and some for more than 10 years.

From before the first session to the end of the last session, the participants improved their [grip strength](#).

"About midway through the sessions, I could open my bedroom door with my left hand for the first time since my [injury](#) and could open new water bottles, when previously someone else had to do this for me," said Cecilia Villarruel, one of the participants, whose injury resulted from a car accident 13 years earlier. "Most people with a [spinal cord](#) injury say they just want to go to the bathroom like a normal person again," she said. "Small accomplishments like opening jars, bottles and doors enable a level of independence and self-reliance that is quite satisfying, and have a profound effect on people's lives."

In addition to regaining use of their fingers, the research subjects also gained other health benefits, including improved blood pressure, bladder function, cardiovascular function and the ability to sit upright without support.

"Within two or three sessions, everyone started showing significant improvements, and kept improving from there," said the study's lead author, UCLA research scientist Parag Gad.

"After just eight sessions, they could do things they haven't been able to do for years," said V. Reggie Edgerton, senior author of the research and a UCLA distinguished professor of integrative biology and physiology, neurobiology and neurosurgery.

This is the largest reported recovery of the use of hands that has been reported in patients with such severe spinal cord injuries, Edgerton said.

The researchers placed electrodes on the skin to stimulate the circuitry of the spinal cord. They call their method "transcutaneous enabling motor control," or tEmc. In the stimulation, electrical current is applied at varying frequencies and intensities to specific locations on the spinal cord.

In the training sessions, the participants squeezed a small gripping device 36 times (18 times with each hand) and held their grip for three seconds; the researchers measured the amount of force they used. The training consisted of two sessions a week over four weeks; the eight sessions each lasted about 90 minutes.

"The combination of spinal stimulation plus training with the hands allows them to regain the lost function," Gad said. They were less dependent on their caregivers, and could feed and dress themselves, he added.

Two of the six returned to Edgerton's laboratory 60 days after the training ended and maintained their grip strength; they could turn a doorknob with one hand, twist off a bottle cap and use a fork with one hand. (The four others did not return to the laboratory. The research subjects live in New York, Minnesota and elsewhere.)

The research is published online this month in the *Journal of Neurotrauma*.

More than 1.2 million Americans are living with paralysis from spinal cord injuries, according to the Christopher and Dana Reeve Foundation. Hundreds of thousands of people in the United States have lost control of vital body functions due to such injuries, including use of their hands and fingers.

"Improved [hand](#) function can mean the difference between needing around-the-clock care and living more independently," said Peter Wilderotter, president and CEO of the Reeve Foundation. "These findings bring great hope to those who were told recovery following paralysis would be impossible. As new discoveries and breakthroughs are uncovered, it is clear the word 'impossible' no longer applies to spinal cord injury."

Edgerton's research team has worked with more than two dozen people with severe [spinal cord injuries](#), the vast majority of whom have shown substantial improvements.

"Nearly everyone thought the only people who would benefit from treatment were those who had been injured for less than a year; that was the dogma. Now we know the dogma is dead," said Edgerton, who is also affiliated with the University of Technology Sydney's Centre for Neuroscience and Regenerative Medicine in Australia. "All of our subjects have been paralyzed for more than a year. We know that in a high percentage of subjects who are severely injured, we can improve their quality of life."

Edgerton is seeking FDA approval for the motor control device so that it can be used by rehabilitation clinics and others. He is able to accept only a small number of people into his research program.

The spinal stimulation approach is inexpensive, does not require surgery and can be used in poor communities and countries without advanced

medical facilities—"and the effects are in some ways, we think, better than surgery," Edgerton said.

"I get criticized a lot for giving 'false hope' but we follow where the science tells us to go and just give the research results," Edgerton said.

"Everything is telling us the nervous system is much more adaptable than we've given it credit for, and can relearn and recover from severe injury."

More information: Parag Gad et al. Noninvasive activation of cervical spinal networks after severe paralysis, *Journal of Neurotrauma* (2018).

[DOI: 10.1089/neu.2017.5461](https://doi.org/10.1089/neu.2017.5461)

Provided by University of California, Los Angeles

Citation: Noninvasive spinal stimulation method enables paralyzed people to regain use of hands, study finds (2018, April 26) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2018-04-noninvasive-spinal-method-enables-paralyzed.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--