

# EULAR recommendations for pain management in inflammatory arthritis and osteoarthritis

May 14 2018

---

The European League Against Rheumatism, EULAR, has published a set of recommendations, designed to help health professionals implement approaches to pain management in inflammatory arthritis and osteoarthritis patients. The aim is to reduce pain and the burden on the individual and society.

Pain is the predominant symptom for people with [inflammatory arthritis](#) (IA) and osteoarthritis (OA), mandating the development of new, evidence-based EULAR Recommendations for the health professional's approach to pain management. The target users of these recommendations are health professionals in the field of rheumatology, including rheumatologists. Prompt and knowledgeable pain management support can reduce pain, increase functioning and well-being, and reduce individual and societal costs. Practitioners in all healthcare settings should therefore have the knowledge and skills required to help people with IA and OA to better manage their pain; rheumatology health professionals are ideally placed to provide comprehensive, evidence-based, and patient-centred care.

Pain management typically includes education. Based on the needs of the patient, education is complemented by physical activity and exercise, aids and assistive devices, psychological and social interventions, sleep hygiene education, weight management, pharmacological and joint-specific treatment options, and interdisciplinary pain [management](#).

Underpinned by available systematic reviews and meta-analyses, these recommendations enable [health professionals](#) to provide knowledgeable [pain management](#) support for people with IA and OA.

A multidisciplinary task force including patient representatives conducted a systematic literature review to evaluate evidence regarding effects on pain of multiple treatment modalities. The task force that included 18 members from 12 countries, consisted of patient representatives, nurses, physiotherapists, psychologists, rheumatologists, a general practitioner, an occupational therapist, a clinical epidemiologist, and a research fellow.

From 2,914 studies initially identified, 186 met inclusion criteria. Based on the reviewed studies and expert opinion, the [task force](#) emphasised the importance for the health professional to adopt a patient-centred framework within a biopsychosocial perspective, to have sufficient knowledge of IA and OA pathogenesis, and to be able to differentiate localised and generalised pain. The assessment of patient needs, preferences and priorities; pain characteristics; previous and ongoing pain treatments; inflammation and joint damage; and psychological and other [pain](#)-related factors were deemed important.

**More information:** EULAR Recommendations for the health professional's approach to pain management in inflammatory arthritis and osteoarthritis: [www.eular.org/sysModules/obxCo...management\\_ia\\_oa.pdf](http://www.eular.org/sysModules/obxCo...management_ia_oa.pdf)

Provided by European League Against Rheumatism (EULAR)

Citation: EULAR recommendations for pain management in inflammatory arthritis and osteoarthritis (2018, May 14) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2018-05-eular-pain-inflammatory-arthritis-osteoarthritis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.