

Some calories more harmful than others

15 May 2018, by Amy Quinton



Credit: CC0 Public Domain

While calories from any food have the potential to increase the risk of obesity and other cardiometabolic diseases, 22 nutrition researchers agree that sugar-sweetened beverages play a unique role in chronic health problems. The disease risk increases even when the beverages are consumed within diets that do not result in weight gain.

It's just one of the conclusions published today in *Obesity Reviews* in a position paper by a group of researchers who participated in the 2017 CrossFit Foundation Academic Conference. The task of researchers was to deliberate the question: Are all calories equal with regards to effects on cardiometabolic [disease](#) and obesity? The paper provides an extensive review of the current science on diets that can lead to obesity, cardiovascular disease and Type 2 diabetes.

The paper's sugar-sweetened beverage consensus is particularly relevant in light of a recent legal battle over warning labels on soda, which hinged on the 9th Circuit Court's determination of whether soda and other sweetened beverages are uniquely harmful to human health or one source of calories among

many.

"What's new is that this is an impressive group of scientists with vast experience in nutrition and metabolism agreeing with the conclusion that [sugar-sweetened beverages](#) increase cardiometabolic risk factors compared to equal amounts of starch," said lead author Kimber Stanhope, a research nutritional biologist with the School of Veterinary Medicine at the University of California, Davis.

Sugar substitute won't make you fat

Another interesting point of consensus among researchers is the role of the sugar substitute aspartame. The authors agreed that aspartame does not promote weight gain in adults. Stanhope said this might come as a surprise to most people.

"If you go on the internet and look up aspartame, the layperson would be convinced that aspartame is going to make them fat, but it's not," said Stanhope. "The long and short of it is that no human studies on noncaloric sweeteners show [weight gain](#)."

The authors also agreed that consumption of polyunsaturated (n-6) fats, such as those found in some vegetable oils, seeds and nuts, lowers disease risk when compared with equal amounts of saturated fats. However, that conclusion comes with a caveat. Dairy foods such as cheese and yogurts, which can be high in saturated fats, have been associated with reduced cardiometabolic risk.

The paper reviews the significant challenges involved in conducting and interpreting nutrition research.

"We have a long way to go to get precise answers on a lot of different nutrition issues," said Stanhope. "Nevertheless, we all agree that a healthy diet pattern consisting of minimally processed whole grains, fruit, vegetables, and healthy fats promotes health compared with the refined and palatable typical Western diet pattern."

More information: K. L. Stanhope et al. Pathways and mechanisms linking dietary components to cardiometabolic disease: thinking beyond calories, *Obesity Reviews* (2018). [DOI: 10.1111/obr.12699](https://doi.org/10.1111/obr.12699)

Provided by UC Davis

APA citation: Some calories more harmful than others (2018, May 15) retrieved 24 May 2018 from <https://medicalxpress.com/news/2018-05-calories.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.