


# Stroke prevention drug combo shows promise, study says

16 May 2018



Micrograph showing cortical pseudolaminar necrosis, a finding seen in strokes on medical imaging and at autopsy. H&E-LFB stain. Credit: Nephron/Wikipedia

If you've had a minor stroke or a transient ischemic stroke (TIA), taking the clot-preventing drug clopidogrel along with aspirin may lower your risk of having a major stroke within the next 90 days, according to new research published in *The New England Journal of Medicine*.

An international study of 4,881 adults in 10 countries who either had a minor stroke or a TIA showed that people who took clopidogrel plus aspirin had a 25 percent lower risk of a major stroke, heart attack or death from blood clots within the three months after the first incident, compared with those who took aspirin alone.

"The study gives us solid evidence that we can use this drug combination to prevent strokes in the highest-risk people, but not without some risk of bleeding," said lead author Clay Johnston, M.D., Ph.D., dean and professor of neurology at Dell Medical School at The University of Texas at Austin.

Both minor stroke and TIA are warning signs that a person has a 3 to 15 percent chance of having a more severe stroke within the next three months. Minor stroke is one that results in mild, nondisabling symptoms. TIA, often called a [warning stroke](#) or mini-stroke, is caused by a temporary blockage in a blood vessel to the brain that often dissolves or dislodges on its own to stop symptoms. More than a third of U.S. adults have had TIA symptoms, according to the American Stroke Association.

The study also showed a small increase in the risk of hemorrhage in the clopidogrel-aspirin group compared with the aspirin alone group. For every 1,000 patients treated with the combination, an extra five major bleeds would be expected but with 15 fewer strokes and other major ischemic events. Because the bleeding events are generally reversible, the overall benefit outweighs the risk for most patients, Johnston said.

"Of the 33 major hemorrhages that occurred in these 4,881 patients, more than half involved the gastrointestinal tract, and none of them was fatal. These largely preventable or treatable bleeding complications of the treatment have to be balanced against the benefit of avoiding disabling strokes," said co-author J. Donald Easton, M.D., professor of neurology at the University of California, San Francisco School of Medicine.

One previous trial in this area showed that clopidogrel plus aspirin was effective in lowering risks but did not find the risk of hemorrhage.

"The results of this large international trial, when added to the results of previous research, provide evidence to support the use of clopidogrel plus aspirin for 90 days among patients with minor ischemic stroke and high-risk TIA treated within 12 hours," said Ralph Sacco, M.D., M.S., professor of neurology at Miller School of Medicine at the University of Miami. "This trial is likely to change

practice since most clinicians and patients are usually willing to accept the increased risk of hemorrhage to offset the disabling impact of a stroke."

The research was part of the Platelet-Oriented Inhibition in New TIA and minor ischemic stroke (POINT) trial—a randomized, double-blind, placebo-controlled trial conducted between May 2010 and December 2017. It included patient information from 269 sites in 10 countries throughout North America, Europe and Australia. Patients were included if they had a minor stroke or a transient [ischemic stroke](#) and were at high risk of having a major stroke.

"It's likely we will see more [patients](#) who have had a TIA or a minor stroke receiving the combination of clopidogrel and aspirin in the future," Johnston said. "If you've suffered from a [minor stroke](#) or TIA, it's important to see a physician immediately, even in the emergency room, to ensure you're taking steps to avoid a potentially debilitating stroke later on," he said. "There are several tests that need to be done right away to determine the cause of the event and to make sure the best treatments are started as soon as possible."

Clopidogrel, known by the brand name Plavix, is a platelet inhibitor commonly prescribed to people who have peripheral artery disease or who have had a recent heart attack or stroke to prevent future cardiac or [stroke](#) events.

Provided by University of Texas at Austin

APA citation: Stroke prevention drug combo shows promise, study says (2018, May 16) retrieved 24 May 2018 from <https://medicalxpress.com/news/2018-05-drug-combo.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*