

# Acupuncture and cognitive behavioral therapy for treating insomnia in cancer survivors

17 May 2018

A Patient-Centered Outcomes Research Institute (PCORI)-supported randomized clinical trial of cancer survivors showed that eight weeks of either acupuncture or cognitive behavioral therapy for insomnia (CBT-I) decreased the severity of insomnia among cancer survivors, though improvements were greatest among patients receiving cognitive behavioral therapy. The study will be presented at the upcoming 2018 ASCO Annual Meeting in Chicago.

"Up to 60 percent of cancer survivors have some form of [insomnia](#), but it is often underdiagnosed and undertreated," said lead study author Jun J. Mao, MD, Chief, Integrative Medicine Service, Memorial Sloan Kettering Cancer Center, New York. "Our trial showed that both CBT-I and acupuncture were effective in treating moderate to severe insomnia, although CBT-I was more effective for those with mild symptoms of insomnia. Now patients have more choices to manage their insomnia."

CBT-I is a newer form of psychotherapy that attempts to modify emotions, behaviors, and thoughts related to sleep. CBT-I has been the gold standard for treatment of insomnia, said Dr. Mao.

To find a therapy to compare with CBT-I, the researchers consulted a group of patient advisors who had cancer and who were knowledgeable about how insomnia could impact their health. Additionally, a survey of cancer survivors found that survivors preferred a natural, non-medicinal approach to treating insomnia. Based on this feedback, and results from other sleep studies that showed it could be beneficial, acupuncture was deemed a reasonable comparison to be used in this trial.

The survivors in the trial had completed cancer

treatment, and the mean time since cancer diagnosis was about six years. The survivors had received treatment for breast, prostate, head and neck, hematologic, and colorectal cancer. In addition, 6 percent had received treatment for more than one type of [cancer](#).

All trial participants had been clinically diagnosed with insomnia by research staff through structured clinical interviews and were randomly assigned to receive either CBT-I or acupuncture (needles placed at a pre-determined set of points on the body shown to influence sleep, pain, and depression) for eight weeks.

The participants who received CBT-I worked with a therapist to re-establish a restorative sleep schedule by:

- Reducing the amount of time spent in bed
- Limiting activities performed in bed to only sleep and sexual activity
- Modifying unhelpful beliefs about sleep
- Promoting good sleep hygiene (avoiding activities that included light from tablets and cellphones, eating too late, and performing vigorous activities; they also set a regular sleep schedule)

Reduction in insomnia severity, measured by the Insomnia Severity Index (ISI), from study entry to week 8 (end of treatment), was the primary study outcome. Survivors were also reassessed 20 weeks after having started the trial. The ISI is a questionnaire that asks people to rate the severity of insomnia problems, such as difficulty falling asleep and staying asleep, and the impact of insomnia on their daily functioning and quality of life. The ISI score ranges from 0-28, with scores 0-7 considered as no clinically significant insomnia, 8-14 mild insomnia, 15-21 moderate insomnia, and

22-28 severe insomnia. At the beginning of the trial, 33 survivors had mild insomnia, 94 had moderate insomnia, and 33 severe insomnia.

## Key Findings

CBT-I was the more effective [treatment](#) overall: After eight weeks, insomnia severity scores fell 10.9 points, from 18.5 to 7.5 for those who received CBT-I vs. 8.3 points for those who received acupuncture treatments, from 17.55 to 9.23. Among people with mild insomnia at the start of the trial, far more had an improvement with CBT-I than with acupuncture (85 percent vs. 18 percent). Participants who started the trial with moderate to severe insomnia had somewhat similar response rates to CBT-I vs. acupuncture (75 percent vs. 66 percent). All survivors maintained improvement in insomnia up to 20 weeks after the start of the trial.

This trial was a comparison between two interventions and determined which approach provided a greater relief of insomnia. Future research will focus on how best to deliver effective treatments to more diverse groups of [cancer survivors](#) to improve sleep management.

Provided by American Society of Clinical  
Oncology

APA citation: Acupuncture and cognitive behavioral therapy for treating insomnia in cancer survivors (2018, May 17) retrieved 17 October 2019 from <https://medicalxpress.com/news/2018-05-acupuncture-cognitive-behavioral-therapy-insomnia.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*