

Stronger alcohol policies help reduce alcohol-related crash deaths in US

29 May 2018



Credit: CC0 Public Domain

Stronger alcohol policies, including those targeting both excessive drinking and driving while impaired by alcohol, reduce the likelihood of alcohol-related motor vehicle crash deaths, according to a new study from Boston Medical Center and Boston University. The findings, published in *JAMA Internal Medicine* and based on national data, demonstrate that policies involving alcohol consumption and drinking and driving contribute to lower rates of alcohol involvement in car crashes, indicating the need for more comprehensive policy approaches to be developed and implemented nationwide.

Alcohol is an established risk factor for car [crash](#) deaths. Approximately 30 percent of deaths come from crashes where one or more drivers had a [blood alcohol concentration](#) (BAC) at or above 0.08 percent, the legal limit for driving in the US. In addition, another 20 percent of deaths come from crashes that involve alcohol at BACs below the current limit. However, little is known about how alcohol policies, which include drinking-related policies (e.g., higher alcohol taxes) and restrictions around drinking and driving (e.g., sobriety

checkpoints) relate to alcohol-related crash deaths.

The new study looked at the relationship between alcohol policies in states and the likelihood of alcohol involvement among those who die in motor vehicle crashes. Data on crash deaths was obtained from the Fatality Analysis Reporting System, while state alcohol policies for each year were characterized using the Alcohol Policy Scale, a measure developed at BMC that looks into the broader alcohol [policy](#) environment based on 29 separate alcohol control policies.

Of all adult crash deaths, stronger, more restrictive state alcohol policies decreased the likelihood of alcohol involvement in crash deaths. Specifically, a one percent increase in the restrictiveness of policies corresponded to a one percent reduction in the likelihood that a crash was alcohol-related (e.g., involved one or more drivers with a BAC at or above the legal limit). Across all states, a 10 percent increase in the restrictiveness of policies would translate into approximately 800 fewer deaths annually.

"Given the risks involved with alcohol use, strengthening alcohol control policies could help prevent many crash deaths, including the 40 percent of deaths that affect victims who are not themselves driving while intoxicated," said Timothy Naimi, MD MPH, the study's lead author who is a physician at BMC's Grayken Center for Addiction and a professor at Boston University Schools of Medicine and Public Health.

Importantly, stronger policies were similarly protective against car crash deaths that involved [alcohol](#) at BACs below the current legal limit. "Although not reflected in our current laws, the risk of crashes starts to increase at BAC levels well below 0.08 percent, so stronger policies offer a way to reduce those deaths as well," said Naimi. At present, most developed nations have BAC limits of 0.05 percent or less, and recently the National

Academies of Sciences and the National Transportation Safety Board have called for lowering permissible BAC limits for driving in the US.

Provided by Boston Medical Center

APA citation: Stronger alcohol policies help reduce alcohol-related crash deaths in US (2018, May 29) retrieved 18 September 2021 from <https://medicalxpress.com/news/2018-05-stronger-alcohol-policies-alcohol-related-deaths.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.