

Prompts may up goals-of-care dialogues at outpatient visits

31 May 2018



documentation, and quality of goals-of-care communication during routine outpatient visits and increased goal-concordant care at three months among patients with stable goals," the authors write.

More information: [Abstract/Full Text Editorial \(subscription or payment may be required\)](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

(HealthDay)—A patient-specific preconversation communication-priming intervention targeting patients and clinicians is associated with an increase in goals-of-care discussions among patients with serious illness, according to a study published online May 26 in *JAMA Internal Medicine*.

J. Randall Curtis, M.D., M.P.H., from the University of Washington in Seattle, and colleagues randomized clinicians to the bilateral, preconversation, communication-priming intervention (65 clinicians; 249 [patients](#)) or usual care (67 clinicians; 288 patients). The trial was conducted among patients with serious illness in outpatient clinics.

The researchers observed a significant increase in goals-of-care discussions at the target visit in the intervention versus the control group (74 versus 31 percent; P

"This [intervention](#) increased the occurrence,

APA citation: Prompts may up goals-of-care dialogues at outpatient visits (2018, May 31) retrieved 22 September 2019 from <https://medicalxpress.com/news/2018-05-prompts-goals-of-care-dialogues-outpatient.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.