

Nurses outperform doctors in colorectal cancer surveillance model

12 June 2018

Putting nurses in charge of colorectal cancer surveillance instead of doctors has shown to reduce the number of unnecessary colonoscopies and the number of cases progressing to cancer.

The South Australian audit of 732 colonoscopies over three months in 2015 found that 97 per cent of procedures adhered to Australian guidelines in the nurse-led model while compliance was just 83 per cent physician-led [surveillance](#) model.

Led by Flinders Medical Centre Senior Research Scientist Dr. Erin Symonds, the research has been published in the *Medical Journal of Australia*.

Colorectal cancer (CRC) is one of the world's most prevalent cancers. Early diagnosis has a major impact on survival and can be achieved by screening with faecal occult blood tests (FOBT) or colonoscopy.

Performing colonoscopies too frequently increases costs and the risk of complications, but delays can increase the risk of CRC.

"A family history of CRC or a personal history of adenoma can increase a person's risk of CRC as much as fourfold, and such individuals are advised to undergo regular surveillance colonoscopy," Dr. Symonds and colleagues based at the based at the Flinders Centre for Innovation in Cancer wrote.

"Screening and surveillance guidelines aim to optimise the effectiveness of CRC prevention, with surveillance intervals generally ranging between one and five years."

"Compliance with those recommendations is poor, however, with as many as 89 per cent of patients receiving inappropriate surveillance, usually a colonoscopy before the recommended date."

Based in the South Australian capital of Adelaide, the Southern Co-operative Program for the

Prevention of Colorectal Cancer (SCOOP) was established in 1999 to improve surveillance rates to match the Australian National Health and Medical Research Council (NHMRC) guidelines on CRC prevention.

SCOOP was expanded to use two models: one nurse-led in public academic hospitals, where nurses make recommendations based on NHMRC guidelines, which are then confirmed by a physician; and the other, physician-led in private non-academic hospitals where the specialist physician manages the entire process.

Following the introduction of the nurse-led model in 2000, adherence to guidelines increased from 46 per cent to 96 per cent within two years and has since been maintained, demonstrating the long-term sustainability of the model.

In the latest research, the nurse-led model of colorectal cancer surveillance improved compliance with guideline recommendations, reduced the number of unnecessary colonoscopies and reduced the number of cases progressing to cancer by a significant number when compared to the physician-led model.

"Having a process in place that allows for long-term compliance with surveillance guidelines will promote optimal health care, as procedures performed too frequently can increase risks to patients, are expensive, and lengthen waiting lists," Dr. Symonds and colleagues wrote.

"Continuous monitoring of and education about colonoscopy surveillance intervals for patients at elevated risk of CRC promotes adherence to recall guidelines and efficient use of limited endoscopy resources."

Dr. Symonds and her colleagues concluded that the nurse-led [model](#) for decisions about colorectal [cancer](#) surveillance intervals achieves a high

degree of compliance with guideline
recommendations, which should relieve burdening
of colonoscopy resources.

Provided by The Lead Australia

APA citation: Nurses outperform doctors in colorectal cancer surveillance model (2018, June 12)
retrieved 17 June 2019 from <https://medicalxpress.com/news/2018-06-nurses-outperform-doctors-colorectal-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.