

CDC: invasive MRSA more likely among injection drug users

12 June 2018



"Continued increases in nonsterile injection drug use are likely to result in increases in invasive MRSA infections, underscoring the importance of public health measures to curb the [opioid epidemic](#)," the authors write.

One author disclosed financial ties to the pharmaceutical industry.

More information: [Abstract/Full Text](#)

Copyright © 2018 [HealthDay](#). All rights reserved.

(HealthDay)—Injection drug users are more than 16-fold more likely to develop invasive methicillin-resistant *Staphylococcus aureus* (MRSA) infections, according to research published in the June 8 issue of the U.S. Centers for Disease Control and Prevention's *Morbidity and Mortality Weekly Report*.

Kelly A. Jackson, M.P.H., from the CDC in Atlanta, and colleagues examined the effect of the opioid epidemic on invasive MRSA infections during 2005 to 2016 using data from the Emerging Infections Program.

The researchers found that the likelihood of developing invasive MRSA infections was 16.3 times higher for persons who inject drugs versus others. From 2011 to 2016 there was an increase in the proportion of invasive MRSA cases that occurred among persons who inject drugs, from 4.1 to 9.2 percent. The [infection](#) types were often those linked to nonsterile injection drug use.

APA citation: CDC: invasive MRSA more likely among injection drug users (2018, June 12) retrieved 20 August 2018 from <https://medicalxpress.com/news/2018-06-cdc-invasive-mrsa-drug-users.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.