

New study links poor sleep quality to atrial fibrillation

26 June 2018



Credit: CC0 Public Domain

Poor sleep quality appears to be an important risk factor for atrial fibrillation, report scientists in the first study of its kind to demonstrate a relationship between poor sleep quality independent of sleep apnea and a higher risk of atrial fibrillation (AF). Their findings are published in *HeartRhythm*.

AF is an irregular, rapid heart rate that may cause symptoms such as heart palpitations, fatigue, and shortness of breath. It can substantially reduce [quality](#) of life and is associated with heightened risks of stroke, dementia, heart attack, kidney disease, and death. Obstructive sleep apnea has been established as a risk factor for AF, but the mechanism is unclear. While episodes of abnormally slow or shallow breathing (hypopnea) and apnea may cause cardiopulmonary stress, induce inflammation, and contribute to cardiovascular disease, [obstructive sleep apnea](#) also results in poor sleep. Aspects of poor sleep such as altered sleep duration, efficiency, and architecture have been linked to other cardiovascular diseases.

has previously been demonstrated, the effect of sleep itself on AF risk has remained unknown," explained lead investigator Gregory M. Marcus, MD, MAS, Electrophysiology Section, Division of Cardiology, Department of Medicine, University of California, San Francisco, CA, USA. "Strategies to enhance [sleep quality](#) are different from those that focus on relieving airway obstruction, so it is important to understand the relationship between sleep itself and AF."

Investigators drew on four different studies to determine whether [poor sleep](#) itself is a risk factor for AF. First, they used the global, internet-based Health eHeart Study and determined that individuals with more frequent nighttime awakenings while trying to sleep more often carried a diagnosis of AF. They then validated these findings by using the NIH-funded Cardiovascular Health Study, a prospective cohort study, in which they found that individuals who reported more frequent nighttime awakenings at baseline exhibited a higher risk of developing AF both before and after adjustment for potential confounders.

Within a subset of these individuals who had undergone formal sleep studies, they found that less REM sleep in particular predicted future AF. Finally, in order to see if these findings were readily translatable to patients already seen in healthcare settings and recognized by their providers as having sleeping difficulty, they drew on the California Healthcare Cost and Utilization Project (HCUP), a set of medical records databases of all California residents aged 21 or older who received care in a California ambulatory surgery unit, emergency department, or inpatient hospital unit between January 2005 and December 2009. Among several million people, the HCUP data confirmed that a diagnosis of insomnia predicted a diagnosis of AF both before and after adjustment for potential confounding effects.

"While a relationship between sleep apnea and AF These results provide more evidence that sleep

quality is important to cardiovascular health and specifically to AF. Investigators determined that there was no evidence that sleep duration per se was a risk factor for AF. Instead, they consistently found [sleep disruption](#) to be an important risk factor. While the underlying mechanisms are still unknown, these findings may motivate novel ways to think about, and hence future research into, factors that influence AF risk.

This is the first study to demonstrate a relationship between worse sleep quality independent of sleep apnea and a higher risk of AF. "These data provide compelling evidence that sleep quality itself, even independent of [sleep apnea](#), is an important determinant of AF risk," noted Dr. Marcus. "While there are several available treatments for AF, prevention of the disease would be ideal. The good news is that sleep quality can be modifiable and is something that at least to some degree is under the control of the individual. It's possible that improving [sleep hygiene](#), such as performing regular exercise, getting to bed at a reasonable hour on a regular basis, and avoiding viewing screens before bed as well as caffeine later in the day, might help stave off AF."

More information: "Sleep characteristics that predict atrial fibrillation," *HeartRhythm*, [DOI: 10.1016/j.hrthm.2018.05.008](#)

Provided by Elsevier

APA citation: New study links poor sleep quality to atrial fibrillation (2018, June 26) retrieved 18 June 2019 from <https://medicalxpress.com/news/2018-06-links-poor-quality-atrial-fibrillation.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.