


Change in brain cells linked to opiate addiction, narcolepsy

June 27 2018



The left image depicts hypocretin-producing neurons in a healthy human brain, on the right, an excess of hypocretin-producing neurons in a heroin user. Credit: UCLA Health



Two discoveries—one in the brains of people with heroin addiction and

the other in the brains of sleepy mice—shed light on chemical messengers in the brain that regulate sleep and addiction, UCLA researchers say.

In 2000, UCLA researchers discovered that human [narcolepsy](#) is caused by a loss of roughly 90 percent of the 80,000 brain cells containing hypocretin, also called orexin, a chemical messenger, or neurotransmitter, important in the regulation of sleep. Narcolepsy causes excessive sleepiness, sleep paralysis (a feeling of being conscious but unable to move), hallucinations and episodes of cataplexy, a partial or total loss of muscle control that is often triggered by a strong emotion such as laughter.

Narcolepsy affects about one in 2,000 to 3,000 people but can go undiagnosed for years, according to the National Sleep Foundation. The sleep disorder usually has its onset in childhood or adolescence.

In a new study, these researchers discovered that people addicted to heroin have, on average, 54 percent more hypocretin-producing neurons than do people without addiction. They confirmed, in mice, that opiates cause this increase. The increase in hypocretin cells lasted for as long as four weeks after discontinuation of the morphine treatment, well after morphine had left the animals' bodies.



Morphine administration restored levels of hypocretin neurons in mice with narcolepsy (bottom), which previously lacked hypocretin neurons when given saline (top). Credit: T.C. Thannickal et al., *Science Translational Medicine* (2018)

The researchers hypothesized that morphine, the active ingredient in heroin, might restore the missing hypocretin-producing neurons in

people with narcolepsy. To test this idea, researchers used mice that were made narcoleptic by the loss of hypocretin cells and gave them morphine. The researchers found that morphine increased the number of hypocretin producing cells and their narcoleptic symptoms disappeared.

The continued production of hypocretin in mice, after discontinuing [morphine](#) treatment, suggests that humans may continue producing hypocretin after they have stopped using heroin. The researchers hypothesize that this hypocretin increase may be a factor in opiate craving. In addition, raising the number of hypocretin cells to near normal levels could potentially reverse narcoleptic symptoms in humans. However, further work in [mice](#) is necessary before this can be recommended as a treatment for narcolepsy in people. Next, researchers would like to find out if reducing "excess" hypocretin [cells](#)—or blocking the action of the hypocretin peptide that they release—could make long-term withdrawal from opiates easier and prevent relapse.

The study appears in the June 27 issue of *Science Translational Medicine*.

More information: T.C. Thannickal et al., "Opiates increase the number of hypocretin-producing cells in human and mouse brain and reverse cataplexy in a mouse model of narcolepsy," *Science Translational Medicine* (2018). [stm.sciencemag.org/lookup/doi/ ... scitranslmed.aao4953](http://stm.sciencemag.org/lookup/doi/10.1126/scitranslmed.aao4953)

Provided by University of California, Los Angeles

Citation: Change in brain cells linked to opiate addiction, narcolepsy (2018, June 27) retrieved 20 September 2024 from <https://medicalxpress.com/news/2018-06-brain-cells-linked-opiate-addiction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.