

Hepatitis C vaccine could dramatically reduce transmission in people who inject drugs

11 July 2018



Equipment used by injecting drug users. Heroin is cooked in the cap above a flame. Cotton is inserted into the heroin inside the cap. Heroin is drawn into the syringe through the cotton. Credit: Larry Ouellet

Among the most serious consequences of the opioid epidemic is the spread of hepatitis C among injecting drug users.

A major new study shows that if a hepatitis C [vaccine](#) were successfully developed, it would dramatically reduce transmission of hepatitis C among [drug](#) users—even though it's unlikely such a vaccine would provide complete immunity.

The study, which employed mathematical modeling, is published in *Science Translational Medicine*.

Four of the study's authors are members of the Program for Experimental and Theoretical Modeling in the division of hepatology of Loyola Medicine and Loyola University Chicago Stritch

School of Medicine. One of the Loyola researchers, Harel Dahari, Ph.D., is a co-senior author of the study, along with Marian Major, Ph.D., of the U.S. Food and Drug Administration. Dr. Dahari is an assistant professor at Loyola University Chicago Stritch School of Medicine.

Vaccines are currently available for hepatitis A and hepatitis B, but a vaccine for hepatitis C is still under investigation. A clinical trial is testing an experimental hepatitis C vaccine on injecting drug users. Unlike many other vaccines, the hepatitis C vaccine is not expected to provide complete immunity, known as sterilizing immunity. A vaccinated person exposed to HCV could still be infected with the virus, although the amount of virus in the bloodstream would be significantly reduced.

The new study calculated how effective a vaccine that provided incomplete immunity would be in preventing transmission among injecting drug users. Researchers developed a mathematical model to determine transmission probabilities in drug users who share needles and syringes. They simulated the sharing of two types of common syringes used by drug users. Using previously published data from people infected or reinfected with hepatitis C virus, researchers then estimated the transmission risks between injecting drug users.




Illustration of hepatitis C transmission via a shared syringe showing how the virus-contaminated blood (red) mixes with the drug (blue) while being injected by an infected individual (A-B). A small amount of the infected mixture remains when a second naïve person uses the same syringe (C-D). Credit: Alexander Gutfriand, Harel Dahari, Marian Major

The study estimated that if an injecting drug user shared a syringe/needle with a second drug user who was infected with hepatitis C, there would be a greater than 90 percent chance the first drug user would also become infected with hepatitis C after six months. However, if a vaccine were used, the transmission risk would decrease to between 1 and 25 percent, depending on the type of needle used and other factors.

"Our findings suggest that a hepatitis C vaccine would be an essential part of a comprehensive prevention strategy to meet the World Health Organization's goal of eradicating hepatitis C by 2030," said study co-author Scott Cotler, MD, head of Loyola's division of hepatology and a professor in the department of medicine of Loyola University Chicago Stritch School of Medicine. Other Loyola co-authors are mathematical modelers Alexander (Sasha) Gutfriand, Ph.D., and Louis Shekhtman, MSc.

Hepatitis C is caused by the hepatitis C virus (HCV). Long-term infection with HCV, known as [chronic hepatitis C](#), usually is silent for many years. But the disease eventually can cause cirrhosis (advanced scarring) of the liver, liver cancer and liver failure. In the United States, as many as 3 million people are chronically infected with HCV, with more than 30,000 new infections per year.

Hepatitis C spreads through contaminated blood, and an estimated 60 percent of HCV infections in the U.S. are attributed to sharing needles, syringes or other drug paraphernalia.

Antiviral drugs are used to treat hepatitis C, with cure rates higher than 90 percent. In addition to stopping the disease from progressing, antivirals also can prevent transmission. However, antivirals are expensive, and many injecting drug users lack access to healthcare in the U.S. And even if they are cured, injecting [drug users](#) can become infected again if they continue to share needles.

"While extremely effective, antivirals alone are unlikely to eliminate hepatitis C globally," Dr. Dahari said. "We need to combine antivirals with a [hepatitis C](#) vaccine and harm-reduction measures such as needle-syringe exchange programs, opioid substitution therapy and behavioral counseling."

Some people think a vaccine needs to be perfect, Dr. Dahari added. "But we found that a vaccine still can be extremely beneficial even if it does not provide complete sterilizing immunity."

More information: M. Major et al., "Modeling of patient virus titers suggests that availability of a vaccine could reduce hepatitis C virus transmission among injecting drug users," *Science Translational Medicine* (2018). stm.sciencemag.org/lookup/doi/...scitranslmed.aao4496

Provided by Loyola University Health System

APA citation: Hepatitis C vaccine could dramatically reduce transmission in people who inject drugs (2018, July 11) retrieved 21 October 2018 from <https://medicalxpress.com/news/2018-07-hepatitis-vaccine-transmission-people-drugs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.