


# Heart attack risk on the rise for pregnant women and death rate remains high

18 July 2018



Credit: CC0 Public Domain

The risk of having a heart attack while pregnant, giving birth, or during the two months after delivery, continues to increase for American women, a new study finds.

As published online July 18 in the *Mayo Clinic Proceedings*, the study, led by NYU School of Medicine researchers, found that the risk of suffering a heart attack among pregnant women rose 25 percent from 2002 to 2014.

The researchers suggest that the trend among many women to have children later in life is one possible reason for the increase, as [heart attack risk](#) rises with age overall, and especially during [pregnancy](#). More women, they say, are also obese and/or have diabetes, key risk factors for heart attack. Another factor that may explain the rising numbers is that myocardial infarcts, the technical name for heart attacks, are easier to detect than a decade ago, as tests for early protein markers of related heart cell damage have improved and become more widely available.

"Our analysis, the largest review in a decade, serves as an important reminder of how stressful

pregnancy can be on the female body and heart, causing a lot of physiological changes, and potentially unmasking risk factors that can lead to heart attack," says study senior investigator and interventional cardiologist Sripal Bangalore, MD, MHA.

As part of the study, researchers examined 49,829,753 births recorded in hospitals, where the majority of deliveries in the United States take place, and found that 1,061 heart attacks happened during labor and delivery. Another 922 women were hospitalized for [myocardial infarction](#) before birth, and 2,390 heart attacks occurred during the recovery period after birth.

Bangalore, an associate professor in the Department of Medicine at NYU Langone Health, says that although the absolute numbers of heart attacks and deaths from them remain low, the persistence of the relatively high death rate (unchanged at 4.5 percent of cases) comes despite advances in treating heart [attacks](#) with drug-coated stents and improved use of blood-thinning medications to prevent heart-vessel blockages.

"Our findings highlight the importance to women considering pregnancy to know their risk factors for heart disease beforehand," says study first author and interventional cardiologist Nathaniel Smilowitz, MD, an assistant professor at NYU Langone. "These patients should work out a plan with their physicians to monitor and control [risk factors](#) during pregnancy so that they can minimize their risk."

The study also provided further evidence that the risk of having a [heart](#) attack during pregnancy rises as women get older. A woman between the ages of 35 to 39 who becomes pregnant is five times more likely to suffer a [heart attack](#) than a woman in her 20s, and women in their early 40s are 10 times more at risk than women in their 20s. Few [women](#), they say, become pregnant after age 45.

Data for the study came from the U.S. Agency for Healthcare Research and Quality's National Inpatient Survey, for which 2014 was the last full year of data available at the time of the analysis. Specifically, rates for myocardial infarction were found to have increased from 7.1 for every 100,000 pregnancies in 2002 to 9.5 for every 100,000 pregnancies in 2014.

Provided by NYU Langone Health

APA citation: Heart attack risk on the rise for pregnant women and death rate remains high (2018, July 18) retrieved 19 October 2020 from <https://medicalxpress.com/news/2018-07-heart-pregnant-women-death-high.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*