

# One in three youth who break the law identify as LGBTQ

1 August 2018

Adolescents who identified as non-heterosexual are significantly over-represented among first-time offenders, according to a new study that examined sexual orientation, gender expression and mental health among adolescents who are involved in the justice system, but are not incarcerated.

The study also found that depression, self-harm, and drug and alcohol use were more prevalent among youth who identified as lesbian, gay, bisexual, trans, queer (LGBTQ) or other sexual minorities, as compared with straight youth.

"It would be incorrect to conclude that sexual-minority status puts a youth at increased risk of offending," said first author Matthew Hirschtritt, MD, MPH, of the UCSF Department of Psychiatry and the UCSF Weill Institute for Neurosciences. "But if we compare court-involved sexual minority youth with court-involved straight youth, we see more severe psychological distress and a greater likelihood of child-welfare system involvement."

More than 70 percent of justice-involved adolescents are in the community and monitored by the courts, rather than in detention, for offenses ranging from underage drinking to breaking and entering. These young people may also be referred to community programs that provide [mental health](#) counseling, substance-use treatment and/or education programs.

Researchers of the UCSF-led study recruited 423 first-time, non-detained offenders, ages 12 to 18, from a juvenile court in a Northeastern state. Of these, 133 adolescents, or 31.4 percent, said they were attracted to or sexually active with their own or both genders, questioned their sexual orientation, or were discriminated against because of their [sexual orientation](#) or gender expression. This percentage is similar to that found in a different study among youth in juvenile detention centers.

Among these 133 sexual-minority adolescents, two reported a different gender identification than that assigned at birth, the authors noted. The study publishes in the *Journal of Adolescent Health* on Aug. 1, 2018.

In contrast, 11.2 percent of U.S. 9th-to-12th graders in a 2015 government-led survey identified as non-heterosexual—just over one-third of the incidence reported in the UCSF-led study.

Hirschtritt and his team found that among the sexual-minority group, whose ages averaged 14, 46.6 percent had used alcohol, versus 26.6 percent of the heterosexual group. They were also approximately twice as likely to use drugs, including synthetic cannabis, cocaine and injection drugs as the heterosexual group—20.2 percent and 10.3 percent respectively.

## **Witnessing Violence, Self-Harm, Substance Abuse Also Common**

The sexual-minority group was more likely to have witnessed violence against a parent or guardian—26.3 percent compared with 11.9 percent for the heterosexual group—and was at higher risk for self-harm, 46.5 percent versus 12.6 percent, respectively.

Additionally, the sexual-minority group reported poorer mental health, as demonstrated by increased difficulties dealing with daily life, inattention and hyperactivity, and internalizing behavior, such as fearfulness and social withdrawal.

There were no significant differences between the two groups' age, race and ethnicity, family income, school-related characteristics or charge types, indicating that sexual-minority status may represent "unique behavioral health risks," Hirschtritt said.

"These results suggest that sexual minority status

must be specifically queried and in multiple ways, so that their needs are met," said senior author Marina Tolou-Shams, Ph.D., of the UCSF Department of Psychiatry, UCSF Weill Institute for Neurosciences and Zuckerberg San Francisco General Hospital and Trauma Center.

"We need cost-effective, tailored interventions so that adolescents' first court appearances become an opportunity to prevent worsening of pre-existing health risks and further entrenchment in the justice system," she added.

"Factors like peer rejection and family discord may contribute to impaired support networks and engagement in risky behavior," said Hirschtritt. "Intake and community supervision staff should query for high-risk behaviors. Additionally, one-on-one support, group counselling and peer support may meet the needs of these adolescents without further stigmatizing them."

Provided by University of California, San Francisco

APA citation: One in three youth who break the law identify as LGBTQ (2018, August 1) retrieved 28 September 2020 from <https://medicalxpress.com/news/2018-08-youth-law-lgbtq.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*