


Function of gene mutations linked to neurological diseases identified

August 10 2018



VPS13 family members mediate lipid transfer between organelles at membrane contact sites. Bottom left: fluorescence image shows the localization of VPS13C (magenta) at contacts between the ER (green) and a lysosome (blue). Bottom right: fluorescence image showing the localization of VPS13A (magenta) at contacts between the ER (green) and a mitochondrion (blue). Credit: Yale University

Several gene mutations have been linked to Parkinson's disease, but exactly how and where some of them cause their damage has been unclear. A new Yale study, published in the *Journal of Cell Biology*, shows that one of the genes whose mutations are responsible for a familial form of Parkinson's encodes a protein that controls the transfer of lipids between membranes of cell organelles.

The labs of Yale's Pietro De Camilli and Karin Reinisch investigated the role of VPS13 proteins whose dysfunction play a role in a variety of neurological disorders, including Parkinson's. They found that VPS13 proteins can extract lipids from cellular membranes, harbor them in a waterproof cavity, and transfer them to adjacent membranes. VPS13 [family members](#) are localized at sites within the cell where organelles are in close proximity to each other—so-called contact sites—and function as tethers between them making lipid transfer more efficient.

VPS13 family members operate at different sites within the cell and are associated with different neurological disorders. VPS13C, the protein associated with Parkinson's disease, acts at contacts between the endoplasmic reticulum (the organelle where most lipids are synthesized) and endosomes/lysosomes (the organelles where most cell components are degraded). Another, VPS13A, is associated with a Huntington's-like syndrome (Chorea-Acanthocytosis) and acts at contacts between the [endoplasmic reticulum](#) and mitochondria, the energy centers of the cell.

As other Parkinson's disease genes also encode proteins of the endo-lysosomal system, the new insight may help identify shared cellular mechanisms in Parkinson's disease resulting from mutations in different genes and thus help design new therapies for this condition, the authors say. Nikit Kumar and Marianna Leonzino of Yale are first authors of the study.

More information: Nikit Kumar et al. VPS13A and VPS13C are lipid

transport proteins differentially localized at ER contact sites, *The Journal of Cell Biology* (2018). [DOI: 10.1083/jcb.201807019](https://doi.org/10.1083/jcb.201807019)

Provided by Yale University

Citation: Function of gene mutations linked to neurological diseases identified (2018, August 10) retrieved 23 April 2024 from <https://medicalxpress.com/news/2018-08-function-gene-mutations-linked-neurological.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.