


Potential biomarker for autism

13 August 2018

Spectrum Disorder, *JNeurosci* (2018). DOI: [10.1523/JNEUROSCI.1229-18.2018](https://doi.org/10.1523/JNEUROSCI.1229-18.2018)



Provided by Society for Neuroscience

Credit: An et al., *JNeurosci* (2018)

A study of young children with autism spectrum disorder published in *JNeurosci* reveals altered brain waves compared to typically developing children during a motor control task. The non-invasive neuroimaging technique used in this study could be employed to detect autism symptoms as early as infancy.

People with autism show reduced inhibitory neural activity, reflected in gamma [brain waves](#), and movement abnormalities.

Mitsuru Kikuchi and colleagues investigated these two aspects of the disorder by recording the brain activity of five- to seven-year-old children diagnosed with autism as they played a video game that required them to press a button to feed a puppy. Compared to a matched group of typically developing children, the children with autism exhibited longer reaction times and reduced gamma oscillations that were associated with the severity of their symptoms.

The researchers demonstrate that these features alone can distinguish the two groups, which may represent an objective approach toward diagnosing autism in [young children](#).

More information: Altered Gamma Oscillations during Motor Control in Children with Autism

APA citation: Potential biomarker for autism (2018, August 13) retrieved 22 September 2019 from <https://medicalxpress.com/news/2018-08-potential-biomarker-autism.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.