

More protein after weight loss may reduce fatty liver disease

August 16 2018

Increasing the amount of protein in the diet may reduce the liver's fat content and lower the risk of diabetes in people with nonalcoholic fatty liver disease (NAFLD). The study is published ahead of print in the *American Journal of Physiology—Endocrinology and Metabolism*.

NAFLD—sometimes referred to as a "fatty [liver](#)"—occurs when more than 5 percent of the liver's total weight is made up of fatty tissue. Excessive fat in the liver can lead to scarring, which may increase the risk of [liver cancer](#) or [liver failure](#). People with NAFLD are more likely to develop type 2 diabetes, and people with type 2 diabetes are more likely to develop NAFLD. In fact an estimated 70 percent of people with type 2 diabetes also have a fatty liver. Obesity is also a major risk factor for NAFLD.

Previous studies have found that short-term protein supplementation helps reduce the fat content in the liver, but there have been few studies on the long-term effects of protein on NAFLD. Researchers conducted a two-year study to determine the long-term impact of [dietary protein](#) on a fatty liver after [weight loss](#). This study was part of the PREVIEW study, which aims to identify the most efficient lifestyle pattern for the prevention of type 2 diabetes in a population of pre-diabetic overweight or obese individuals.

Twenty-five adult volunteers—15 of whom had been previously diagnosed with NAFLD—participated in a low-calorie diet for eight weeks to lose up to 8 percent of their body weight. After weight loss, the

volunteers were directed to maintain their weight for two years and to follow either a moderate- or high-protein diet averaging from 0.8 to 1 grams of protein per kilogram (2.2 pounds) of body weight. The research team took blood and urine samples and performed body scans to assess liver fat content and the amount of protein eliminated from the volunteers' bodies at three intervals: the start of the weight maintenance phase and again six months and then two years later.

After two years maintaining their weight loss, the increase in dietary protein was associated with reduced liver fat content in the volunteers. In addition, more than half of the participants who were previously diagnosed with NAFLD no longer had a fatty liver.

"These findings stress the clinical implications and potential benefits of increased [protein](#) intake after [weight](#) loss for people with NAFLD at risk to develop diabetes," the researchers wrote.

More information: Mathijs Drummen et al, Long-term effects of increased protein intake after weight loss on intrahepatic lipid content and implications for insulin sensitivity - a PREVIEW study, *American Journal of Physiology-Endocrinology and Metabolism* (2018). [DOI: 10.1152/ajpendo.00162.2018](#)

Provided by American Physiological Society

Citation: More protein after weight loss may reduce fatty liver disease (2018, August 16) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-08-protein-weight-loss-fatty-liver.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
