













Assessment of ED threat perceptions identifies patients at risk for cardiac-induced PTSD


2 October 2018

Measuring patients' threat perceptions in ED
 1000 adults evaluated for ACS in ED (English/Spanish speaking)
 12 items tested from meta-analysis

| Included in measure | Not included |
|--|---|
|  I feel vulnerable (FL 0.669) |  I am worried that the doctors are not in control of my situation (FL 0.457) |
|  I am worried that I am not in control (FL 0.665) |  I am concerned about my family (FL 0.454) |
|  I am worried that my symptoms are severe (FL 0.665) |  I believe that my symptoms are connected to my heart (FL 0.381) |
|  I feel helpless (FL 0.628) |  I am in pain (FL 0.267) |
|  I am worried that I am going to die (FL 0.617) |  I feel like I can handle the symptoms myself (FL -0.133) |
|  I am afraid (FL 0.579) | |
|  I think this event will have a big impact on my life (FL 0.563) | |

Cronbach's alpha at bedside 0.82
 FL: factor loading

Cornelius doi 10.1111/acem.13513



seven-item measure of ED threat perceptions (e.g., patients' subjective feelings of helplessness or lack of control) in English- and Spanish-speaking patients evaluated for [acute coronary syndrome \(ACS\)](#).

In addition to providing a foundation for standardized research programs across protocols and disease contexts, the study suggested that the brief assessment could inform clinical interventions and has practical implications: the ability to identify at-risk patients at an early point of contact in the ED and provide appropriate and targeted support to improve patient outcomes. Further, the study showed the feasibility of administering this instrument in the acute care setting.

1,000 adults were evaluated for acute coronary syndrome (ACS) in the emergency department; 12 items were tested from meta-analysis. Credit: Kirsty Challen, Lancashire Teaching Hospitals, United Kingdom

Rosemarie Ramos, Ph.D., MPH, assistant professor and assistant research program director in the Department of Emergency Medicine, University of Texas Health-San Antonio, commented:

A brief tool assessing emergency department (ED) threat perceptions has clinical utility for providers to identify patients at risk for developing cardiac-induced PTSD and is critical to inform research on whether threat may be modified in-ED to reduce post-traumatic stress disorder (PTSD) incidence. That is the finding of a study to be published in the October 2018 issue of *Academic Emergency Medicine* (AEM), a journal of the Society for Academic Emergency Medicine (SAEM).

"Perceived stress and powerlessness are well-established risk factors for poor functional and psychological resilience among those who have witnessed trauma (e.g., community violence, military combat.) When one experiences the trauma of [emergency](#) treatment for a life-altering condition in a seemingly-chaotic setting, the psychological impact of such is rarely addressed at point-of-care. This study offers an invaluable instrument that aligns with the goals of Social Emergency Medicine; specifically, the need to assess and address plausible determinants of resilience before discharge from the ED."

The lead author of the study is Talea Cornelius, Ph.D., a postdoctoral research scientist at Columbia University Medical Center in the Center for Behavioral Cardiovascular Health, New York.

The findings of the study are discussed in the most recent AEM podcast, [Threat Perceptions in the ED](#).

The study, by Cornelius, et al., reports the development and validation of a brief, self-report,

More information: Talea Cornelius et al, Development and Validation of a Measure to Assess Patients' Threat Perceptions in the Emergency Department, *Academic Emergency Medicine* (2018). [DOI: 10.1111/acem.13513](https://doi.org/10.1111/acem.13513)

Provided by Society for Academic Emergency
Medicine

APA citation: Assessment of ED threat perceptions identifies patients at risk for cardiac-induced PTSD (2018, October 2) retrieved 18 October 2019 from <https://medicalxpress.com/news/2018-10-ed-threat-perceptions-patients-cardiac-induced.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.