


Inflammation in the womb may explain why some babies are more prone to sepsis after birth

October 9 2018



Analyses of chorioamnion, lungs, gut, and plasma samples. (A-C) Images from hematoxylin and eosin staining in the chorioamnion, scale bars represent 100 μm . (D-E, G-H) Myeloperoxidase (MPO) staining in lungs and gut, scale bars represent 200 μm . (F) Lung CXCL8 levels. (I-K) Incidence of severe diarrhea and physical activities in postnatal control and LPS-exposed pigs. (L) Plasma interleukin (IL)-1 β levels in the two treatment groups. CON, control pigs; LPS, intra-amniotic lipopolysaccharide exposed pigs. *P *American Journal of Pathology*

Each year 15 million infants are born preterm and face high risks of short- and long-term complications, including sepsis, severe inflammation of the gut, and neurodevelopmental disorders. A new report in the *American Journal of Pathology* demonstrates a link between prenatal inflammation and postnatal immune status and organ function in preterm pigs, suggesting that early intervention (eg, antibiotics or anti-inflammatory drugs) may be warranted for infants born preterm with signs of inflammation of fetal membranes.

"Our study may urge clinicians to be more aware of the population of preterm infants with chorioamnionitis ([inflammation](#) of the fetal membrane) as they have higher risks of systemic inflammation and neonatal sepsis," explained lead investigator Per T. Sangild, DVSc, DMSc, Ph.D., of the Section of Comparative Pediatrics and Nutrition, Department of Veterinary and Animal Sciences, University of Copenhagen, Frederiksberg, Denmark, and the Department of Pediatrics and Adolescent Medicine, Rigshospitalet, Copenhagen, Denmark.

"The data imply the importance of the integrity of barriers between epithelial tissues (eg, gut, lungs, and skin) and the circulation. Those barriers are more fragile in preterm neonates, and they can facilitate

translocation of bacteria and inflammatory molecules, leading to systemic inflammation and internal organ disorders."

To induce prenatal inflammation, the bacterial endotoxin lipopolysaccharide (LPS) was injected into the amniotic sacs of prenatal pigs. LPS and control groups were analyzed at birth, three days after birth, and five days after birth (formula feeding).

At birth, prenatal LPS induced mild histologic chorioamnionitis and strong fetal lung and gut innate immune responses with elevated inflammatory cytokines and neutrophil/macrophage infiltration. "We believe the epithelial responses were likely derived from direct exposure to intra-amniotic LPS or LPS-induced cytokines," noted Dr. Sangild.

Five days later, the gut and [lung inflammation](#) subsided; however, the pigs exposed to LPS prenatally gradually developed systemic inflammation, with high levels of blood leukocyte subsets (eg, neutrophils, lymphocytes) and plasma cytokines (eg, IL-1 β), similar to symptoms found in septic infants. Among those who survived, the pigs in the control group were on their feet and walked for the first time before the LPS-exposed animals. High levels of bacteria were also found in the spleen in the LPS-exposed pigs, indicating increased systemic infection or decreased capacity to clear translocated bacteria. The in utero death rate was higher in the LPS-exposed fetuses compared to the control group, as was the incidence of severe diarrhea. Interestingly, intra-amniotic LPS did not increase the incidence of formula-induced necrotizing enterocolitis (NEC) on Day 5. This is important as it has been suspected that [systemic inflammation](#) at birth might also predispose to later NEC in the gut of [preterm infants](#); however, this hypothesis was not supported by this study.

"These data suggest that the postnatal systemic effects of short-term prenatal LPS were indirectly initiated from the preceding local

inflammation of [epithelial tissues](#) in the fetal period, and the effects were gradually amplified systemically during the first few days after preterm birth," commented Dr. Sangild.

The results of this study highlight the importance of early diagnosis of prenatal inflammation to facilitate nutritional, medical, or pharmaceutical interventions that attenuate the detrimental postnatal responses to prenatal inflammation. The problem still remains that a pregnant woman with intra-amniotic inflammation may be asymptomatic and, therefore, unaware she has an infection that could harm her baby.

More information: "Prenatal Intra-Amniotic Endotoxin Induces Fetal Gut and Lung Immune Responses and Postnatal Systemic Inflammation in Preterm Pigs," *American Journal of Pathology* (2018). [DOI: 10.1016/j.ajpath.2018.07.020](#)

Provided by Elsevier

Citation: Inflammation in the womb may explain why some babies are more prone to sepsis after birth (2018, October 9) retrieved 19 September 2024 from <https://medicalxpress.com/news/2018-10-inflammation-womb-babies-prone-sepsis.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
