

In black men, baseline screening in midlife strongly predicts aggressive prostate cancer

10 October 2018

Black men are more likely to be diagnosed with, and to die from, prostate cancer yet recommendations about prostate cancer screening are primarily based on studies of white men. A new study led by investigators at Brigham and Women's Hospital and Harvard T.H. Chan School of Public Health is the first to look at results exclusively among black men and to address whether an optimized screening strategy with baseline prostate specific antigen (PSA) levels predict prostate cancer in this population. The study finds that baseline PSA levels measured at midlife strongly predicted risk of total and aggressive prostate cancer in black men years in the future. Results are published online in *European Urology*.

"We found that a single, baseline PSA level measured during midlife strongly predicted subsequent diagnosis of total and aggressive [prostate cancer](#) up to 12 years after a blood draw," said co-lead author Mark Preston, MD, MPH, a urologic surgeon in the Division of Urology at BWH. "Our findings suggest that targeted screening based on a midlife PSA might identify black men at high risk of aggressive prostate cancer while minimizing screening in those at low risk."

"Black men in the U.S. are 2.5 times more likely to die of prostate cancer compared to white men, yet screening studies to date have largely been focused on white men," said co-lead author Lorelei Mucci, MPH, ScD, associate professor of Epidemiology at the Harvard T.H. Chan School of Public Health. "As such, the data from our study address a critical gap in understanding the substantial racial disparities and suggests a potential strategy to reduce prostate cancer death in black men."

The research team took a targeted, risk-stratified approach, leveraging data on men enrolled in the federally funded Southern Community Cohort Study (SCCS), a collection of 86,000 men and women from the Southeastern U.S., that included more than 22,000 [black men](#). The team performed a nested, case-control study of 197 men, ages 40 to 64, who developed prostate cancer, including 91 cases of aggressive disease, and compared them to 569 matched controls.

The team found that 95 percent of total and 97 percent of aggressive prostate cancer cases had baseline PSA above the average for their age group. Compared to men with PSA at or below the age-specific median, men with levels above the median had significantly increased risk of aggressive [prostate cancer](#) across age groups. Men with PSA levels above the 90th percentile had the greatest risk.

The authors noted that increased risk was seen with PSA levels that were higher than the average but still well within the "normal" range—and low enough not to trigger follow-up in usual clinical practice.

"These findings do not imply that prostate biopsy or definitive treatment is immediately required in younger men with higher PSA levels at baseline, as this could lead to over-diagnosis, but that they undergo more intensive PSA [screening](#) to enable earlier identification of cancer and potential cure while still possible," the authors write.

Provided by Brigham and Women's Hospital

APA citation: In black men, baseline screening in midlife strongly predicts aggressive prostate cancer (2018, October 10) retrieved 19 June 2021 from <https://medicalxpress.com/news/2018-10-black-men-baseline-screening-midlife.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.