

Concomitant use of sleeping pills and strong painkillers is common among people with Alzheimer's disease

19 November 2018



Credit: CC0 Public Domain

One in five people with Alzheimer's disease (AD) who use a benzodiazepine are also concomitant users of an opioid, according to a new study from the University of Eastern Finland. Concomitant use was more common in comparison persons, but those with AD used strong opioids more frequently. About half of all concomitant users were prolonged users whose use of these drugs had continued for more than three consecutive months.

Prolonged concomitant use was associated with lung diseases, osteoporosis and previous hip fracture. In addition, a history of substance abuse and previous long-term use of [benzodiazepines](#) were related to prolonged concomitant use. In these high-risk patient groups, concomitant use of opioids and benzodiazepines or related drugs may increase the risk of pneumonia, fractures and [drug](#) misuse.

The most commonly used drug combination among persons with AD was a benzodiazepine and a weak opioid, while those without AD most commonly used a benzodiazepine related Z-drug

and a weak opioid. Those with AD were more commonly prescribed strong opioids than those without AD. The trend may be partly explained by the use of [opioid](#) patches that are more commonly used in individuals with AD. In Finland, only buprenorphine and fentanyl are available as patches.

"Concomitant use of drugs that act on the central nervous system in an older population is concerning, as the use of these drugs has been associated with serious risks, especially in frail individuals with AD. Unnecessary co-use of these drugs should be avoided, as the benefits rarely outweigh the risks," says University Lecturer Niina Karttunen from the University of Eastern Finland.

The study was part of the nationwide register-based MEDALZ study that includes all persons with clinically verified diagnosis of Alzheimer's disease in Finland during 2005–2011. The study included 70,718 persons diagnosed with Alzheimer's disease. Data on drug use was extracted from the Finnish Prescription Register. The study was conducted at the University of Eastern Finland and published in the *International Journal of Geriatric Psychiatry*.

More information: Niina Karttunen et al. Concomitant use of benzodiazepines and opioids in community-dwelling older people with or without Alzheimer's disease - A nationwide register-based study in Finland, *International Journal of Geriatric Psychiatry* (2018). [DOI: 10.1002/gps.5018](https://doi.org/10.1002/gps.5018)

Provided by University of Eastern Finland

APA citation: Concomitant use of sleeping pills and strong painkillers is common among people with Alzheimer's disease (2018, November 19) retrieved 24 June 2019 from <https://medicalxpress.com/news/2018-11-concomitant-pills-strong-painkillers-common.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.