

Depressed people have an increased risk of atrial fibrillation

20 November 2018

Depressed people have an increased risk of atrial fibrillation, according to a study published today in the *European Journal of Preventive Cardiology*, a European Society of Cardiology (ESC) journal. Medication was not responsible for the high frequency of atrial fibrillation in depressed people. The findings are reported during Global AF Aware Week.

Study author Morten Fenger-Grøn, senior statistician, Research Unit for General Practice, Aarhus University, Denmark, said: "It is common knowledge that there is a connection between the mind and the heart. Depression predicts the development of coronary artery disease and worsens its prognosis. Our study investigated whether [depression](#) is also linked with atrial [fibrillation](#)."

Atrial fibrillation is the most common heart rhythm disorder (arrhythmia). It causes 20-30% of all strokes and increases the [risk](#) of dying prematurely. One in four middle-aged adults in Europe and the US will develop atrial fibrillation. It is estimated that by 2030 there will be 14-17 million patients with atrial fibrillation in the European Union, with 120,000-215,000 new diagnoses each year. Signs of atrial fibrillation include palpitations, shortness of breath, tiredness, chest pain and dizziness.

Previous studies have found associations between depression and both more severe symptoms and higher mortality in atrial fibrillation patients.⁵ Antidepressants have been linked with some serious, but rare, heart rhythm disturbances, prompting the question of whether they might also raise the risk of atrial fibrillation.

This study investigated the association of depression, and [antidepressant treatment](#), with the risk of developing atrial fibrillation. Filling a prescription for antidepressants for the first time was used as an indicator of depression.

The study included all 785,254 Danish citizens initiating antidepressant treatment from 2000 to 2013 and a 1:5 random sample of the Danish population matched on sex and birth month. The risk of atrial fibrillation was assessed after starting treatment and in the month before, when it was assumed that patients were depressed but medically untreated.

Compared with the general population, patients taking antidepressants had a 3.18-fold higher risk of atrial fibrillation during the first month of treatment. The association gradually reduced thereafter, to 1.37-fold at 2-6 months, and 1.11-fold at 6-12 months.

Mr Fenger-Grøn said: "Filling a prescription for antidepressants, which we used as an indicator of depression, was associated with a three-fold greater risk of atrial fibrillation. The decrease with time could suggest that treatment may alleviate this risk."

The risk of atrial fibrillation risk was even higher in the month before starting antidepressants (7.65-fold). "This suggests that antidepressant medication itself is not associated with the development of atrial fibrillation," said Mr Fenger-Grøn. "If you are depressed, there is no reason to worry that taking drug treatment will cause atrial fibrillation."

Mr Fenger-Grøn noted that in the month before starting medication, the risk of atrial fibrillation was 7.65-fold from 30 to 15 days before, and 4.29-fold in the last 15 days before. He said: "In Denmark it is not typical for doctors to prescribe antidepressants at the first appointment. Our findings indicate that the initial conversation with a doctor may start to ease depressive symptoms."

He added: "The message for patients who already have atrial fibrillation is that you do not need to be concerned about taking antidepressant medication

if you need it. Look after your mental health because our study supports existing evidence that problems with the mind can be detrimental for the heart."

More information: Morten Fenger-Grøn et al. Depression, antidepressants, and the risk of non-valvular atrial fibrillation: A nationwide Danish matched cohort study, *European Journal of Preventive Cardiology* (2018). DOI: [10.1177/2047487318811184](https://doi.org/10.1177/2047487318811184)

Provided by European Society of Cardiology

APA citation: Depressed people have an increased risk of atrial fibrillation (2018, November 20) retrieved 14 November 2019 from <https://medicalxpress.com/news/2018-11-depressed-people-atrial-fibrillation.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.