

# Rerouting nerves during amputation reduces phantom limb pain before it starts

27 December 2018



Dr. J. Byers Bowen (left) and Dr. Ian Valerio follow up with a leg amputee who had targeted muscle reinnervation surgery at The Ohio State University Wexner Medical Center. Those who received the procedure at the time of amputation reported significantly less pain caused by severed nerves. Credit: The Ohio State University Wexner Medical Center

Doctors at The Ohio State University Wexner Medical Center and College of Medicine are pioneering the use of primary targeted muscle reinnervation (TMR) to prevent or reduce debilitating phantom limb and stump pain in amputees.

Losing a limb due to trauma, cancer, or poor circulation can result in phantom limb and stump pain in upwards of 75 percent of amputees in the United States. Primary TMR—the rerouting of nerves cut during [amputation](#) into surrounding muscle—greatly reduces phantom limb and residual limb pain, as reported in recent publications by Dr. Ian Valerio, division chief of Burn, Wound and Trauma in Ohio State's Department of *Plastic and Reconstructive Surgery*, and Dr. J. Byers Bowen, a former resident who is now in private practice. Their latest work featured in the January 2019 issue of *Plastic and Reconstructive Surgery*

describes how to perform this technique in below-the-knee amputations.

TMR was first developed to allow amputees better control of upper limb prosthetics. Traditionally doctors perform the surgery months or years after the initial amputation. When surgeons discovered the procedure also improves certain causes of pain, they started using it to treat disorganized [nerve endings](#) called symptomatic neuromas and/or [phantom limb](#) pain.

In this paper, Valerio and Bowen provide a detailed description of TMR in below-the-knee amputees and document the benefits of primary TMR for preventing pain.

"This paper provides a blueprint for improving [patient outcomes](#) and quality of life following amputation," said Dr. K. Craig Kent, dean of The Ohio State University College of Medicine.

Over the course of three years, the surgeons performed 22 TMR surgeries on below-the-knee amputees, 18 primary and four secondary. None of the patients have developed symptomatic neuromas and only 13 percent of patients who received primary TMR reported having pain six months later.

"A significant amount of pain in amputees is caused by disorganized nerve endings, i.e. symptomatic neuromas, in the residual limb. They form when nerves are severed and not addressed, thus they have nowhere to go," Valerio said. "Attaching those cut nerve endings to motor nerves in a nearby muscle allows the body to re-establish its neural circuitry. This alleviates phantom and residual limb pain by giving those severed nerves somewhere to go and something to do."

Valerio said patients who've had TMR significantly reduce or sometimes stop using narcotics and other nerve pain related medications, which can

greatly improve their quality of life.

minimal health risk and recovery is similar to that of traditional amputation surgery.



Surgeons perform TMR routinely at Ohio State, with primary TMR as the standard of care for most orthopedic-based traumatic and oncologic amputations. Valerio lectures and trains surgeons around the world on the primary TMR technique in an effort to make it a global best practice.

Provided by Ohio State University Medical Center

Robert Haas experienced debilitating pain following the amputation of his left leg that prevented him from using his prosthetic and required medication to manage. However, a surgery that reconnects severed nerves eliminated his pain and allowed him to adjust to life with an artificial limb. Credit: The Ohio State University Wexner Medical Center

"TMR has been shown to reduce pain scores and multiple types of pain via a variety of validated pain surveys. These findings are the first to show that surgery can greatly reduce phantom and other types of limb pain directly," Valerio said.

Bowen added that upper extremity amputees are better able to use and control their prosthetics in addition to their improved pain outcomes. He said, "TMR allows for more individual muscle unit firings through the patient's thoughts. It provides for better intuitive control resulting in more refined functional movements and more degrees of motion by an advanced prosthetic."

The researchers believe primary TMR is a reliable technique to prevent the development of disorganized [nerve](#) endings and to reduce phantom and other [limb pain](#) in all types of amputations. When done at the time of initial amputation, there is

APA citation: Rerouting nerves during amputation reduces phantom limb pain before it starts (2018, December 27) retrieved 23 April 2021 from <https://medicalxpress.com/news/2018-12-rerouting-nerves-amputation-phantom-limb.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*