

# Bile acids mediate metabolic benefits of weight-loss surgery

11 January 2019, by Leigh Macmillan



Vance Albaugh, MD, PhD, left, Robb Flynn, PhD, and Naji Abumrad, MD, seated, are studying how weight-loss operations cause positive metabolic effects, even before weight loss happens. Credit: John Russell

A team of Vanderbilt investigators has pinpointed the role of bile acids and a specific signaling pathway in the positive metabolic effects of weight-loss surgery.

The findings, reported in the journal *Gastroenterology*, also suggest that the intestinal microbiome participates in post-surgery improvements. The results could guide new treatment strategies for obesity and diabetes, which continue to climb in the American population.

Bariatric surgery—weight-loss operations that include [gastric bypass](#)—is much more effective than intensive medical therapy for obesity and diabetes, and metabolic benefits can happen within days, before weight loss occurs.

"Surgical patients can experience dramatic changes in glucose regulation while they are still in the recovery room," said Charles (Robb) Flynn, Ph.D., associate professor of Surgery. "These operations cause metabolic changes that we don't

really understand."

In previous studies, Flynn, Naji Abumrad, MD, John L. Sawyers Professor of Surgical Sciences, and others found increased levels of serum [bile acids](#), gut compounds that help digest dietary fats, after two common bariatric operations—Roux-en-Y gastric bypass and vertical sleeve gastrectomy.

Roux-en-Y gastric bypass, which is one of the most commonly performed weight-loss procedures, restricts the size of the stomach, alters the path of food through the digestive tract and changes the point where bile acids enter the small intestine, from the usual upper part of the small intestine to a site near the end.

Flynn and Abumrad worked with the Vanderbilt Mouse Metabolic Phenotyping Center to develop a simpler surgical procedure in mice that mimics the Roux-en-Y re-routing of bile acids. The procedure, called bile diversion, increases bile acids in the blood and produces all of the beneficial effects of [bariatric surgery](#): weight loss, reduced food intake and improved oral glucose tolerance, Flynn said.

In addition to aiding fat absorption and digestion, bile acids activate signaling pathways through two types of receptors: Tgr5 and Fxr.

In studies led by research fellow Vance Albaugh, MD, Ph.D., the investigators discovered that in mice missing Tgr5, bile diversion surgery still produced beneficial metabolic effects. In mice missing intestinal Fxr, however, bile diversion surgery no longer improved oral glucose tolerance.

"Our study shows specifically that elevated bile acids following bariatric surgery work through the Fxr receptor, and that it functions in the intestine," Flynn said.

The team further found that improvements in glucose handling were accompanied by increased

levels of both GLP-1 (glucagon-like peptide-1) and the intestinal bacteria *Akkermansia muciniphila*, and that the GLP-1 receptor was required for the beneficial effects of bariatric surgery. GLP-1 is a hormone that participates in regulating blood glucose.

"Our mouse model of bariatric surgery has allowed us to isolate the metabolic effects of bile acids and show that improvements in oral glucose handling require an Fxr-GLP-1 receptor signaling pathway," Flynn said.

The change in one species of intestinal bacteria is also intriguing, Flynn said. *Akkermansia muciniphila* is associated with lean phenotypes in animal models and human beings.

"Our discovery raises the idea that finding a way to introduce this bacterium into people might have [weight loss](#) and beneficial metabolic effects," Flynn said. "There's a very tight interplay between bacteria in the gut, bile acids and metabolic effects. We're just beginning to tease this apart."

Flynn and his colleagues are continuing to explore metabolic changes following bariatric surgery in their surgical mouse model. They have also begun to study patients at the Vanderbilt Weight Loss Center to explore whether the mechanisms they've discovered in mice also hold true in people.

"We want to fully understand how the [surgery](#) is generating beneficial metabolic changes so that we can improve treatments for obesity and diabetes," Flynn said.

**More information:** Vance L. Albaugh et al. Role of Bile Acids and GLP-1 in Mediating the Metabolic Improvements of Bariatric Surgery, *Gastroenterology* (2018). [DOI: 10.1053/j.gastro.2018.11.017](#)

Provided by Vanderbilt University

APA citation: Bile acids mediate metabolic benefits of weight-loss surgery (2019, January 11) retrieved 16 January 2019 from <https://medicalxpress.com/news/2019-01-bile-acids-metabolic-benefits-weight-loss.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*