

# Addressing expected challenges after resumption of HPV vaccination

January 17 2019

		Public Subsidies (Age: 13-16)			National Routine Immunization Program (Age: 12-16)						Vac Rate*	
		2010	2011	2012	Suspension of Recommendation			Resumption of Recommendation (hypothesized)				
		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	
Pre-Introduction Generation	1990	20	21	22	23	24	25	26	27	28	29	-
	1991	19	20	21	22	23	24	25	26	27	28	-
	1992	18	19	20	21	22	23	24	25	26	27	-
	1993	17	18	19	20	21	22	23	24	25	26	-
Vaccinated Generation	1994	16	17	18	19	20	21	22	23	24	25	70%
	1995	15	16	17	18	19	20	21	22	23	24	70%
	1996	14	15	16	17	18	19	20	21	22	23	70%
	1997	13	14	15	<del>16</del>	17	18	19	20	21	22	70%
	1998		13	14	<del>15</del>	<del>16</del>	17	18	19	20	21	70%
	1999			13	<del>14</del>	<del>15</del>	<del>16</del>	17	18	19	20	70%
Unvaccinated Generation	2000				<del>13</del>	<del>14</del>	<del>15</del>	<del>16</del>	17	18	19	4%
	2001				<del>12</del>	<del>13</del>	<del>14</del>	<del>15</del>	<del>16</del>	17	18	1%
	2002					<del>12</del>	<del>13</del>	<del>14</del>	<del>15</del>	<del>16</del>	17	0%
	2003						<del>12</del>	<del>13</del>	<del>14</del>	<del>15</del>	<del>16</del>	?
Resumption Generation (hypothesized)	2004						<del>12</del>	<del>13</del>	<del>14</del>	<del>15</del>	<del>16</del>	?

HPV vaccination per birth year. Public subsidies for HPV vaccine started in 2010 in Japan. Nationally, around 70% of targeted girls, aged 13-16, were immunized. HPV vaccination became a national routine immunization for girls aged 12-16 in April of 2013. However, news media repeatedly reported on patients with widespread pain or movement disorders, occurring after immunization. In June of the same year, MHLW announced the suspension of its

recommendation of routine HPV immunization. ©: Around 50-70% of girls have been immunized. ☐: Around 0-20% of girls have been immunized. X: Around 0-10% of girls have been immunized. XX: Vaccination rate was almost 0. △: Around 10-20% will be expected at the highest if the governmental recommendation is resumed in 2019. Credit: Osaka University

In Japan, an HPV vaccination program started in 2010, and the HPV vaccine became a nationally recommended routine immunization for girls aged 12–16 years in April 2013. However, because cases of young girls with widespread pain and movement disorders after vaccination were reported in the media in June 2013, Japan's Ministry of Health, Labour and Welfare (MHLW) announced the suspension of its recommendation for routine HPV immunization.

Previous research at Osaka University showed that the HPV vaccination hiatus would increase the risk of HPV infection and future cervical cancer for girls who did not get vaccinated. HPV vaccination is vital to reduce the risk of HPV infection and cervical cancer; however, no reports on countermeasures against expected challenges after resumption of HPV vaccination have been published.

Researchers at Osaka University compiled countermeasures against predictable challenges after resumption of HPV vaccination, published in *The Lancet Oncology*.

Every year, about 9,000 women are diagnosed with cervical cancer and 2,000 to 3,000 women die of the cancer. A major factor for the development of cervical cancer is infection with HPV, which is mainly transmitted through sexual contact.

This research group demonstrated two possible challenges after

resumption of the MHLW's recommendation for HPV vaccination:

1. Reducing the risk of cervical cancer that will increase by the suspension of the recommendation
2. Promoting HPV vaccination

The group discussed countermeasures against these challenges, making the following proposals for providing information and fostering public acceptance of the vaccine.

1. Easy access to immunization for women who are older than the normally targeted ages of 12–16 years and who were not vaccinated during the suspension of recommendation for HPV vaccination
2. Introduction of the nine-valent vaccine, which can prevent 80-90% of cervical cancer
3. Immunization for boys of the same ages as the targeted girls
4. Reducing health damage due to the suspension of recommendation for HPV vaccination by encouraging medical check-ups and cervical cancer screening
5. Promoting HPV vaccination again by using a behavioral economics approach
6. Providing media with correct information about the HPV vaccine

Dr. Yutaka Ueda says, "Resumption of the government recommendation for HPV vaccination will be insufficient to deal with the expected challenges. It's necessary to reduce negative effects of the suspension of recommendation for HPV vaccination. We hope our proposals will reduce the development of [cervical cancer](#) in Japanese women, and, moreover, protect women's health."

**More information:** Yutaka Ueda et al. Beyond resumption of the

Japanese Government's recommendation of the HPV vaccine, *The Lancet Oncology* (2018). [DOI: 10.1016/S1470-2045\(18\)30573-4](https://doi.org/10.1016/S1470-2045(18)30573-4)

Provided by Osaka University

Citation: Addressing expected challenges after resumption of HPV vaccination (2019, January 17) retrieved 16 April 2024 from <https://medicalxpress.com/news/2019-01-resumption-hpv-vaccination.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.