

Prevalence of knee OA lower with menopausal hormone therapy

17 January 2019



undertaken to establish a causal association between knee OA and MHT," the authors write.

More information: [Abstract/Full Text](#)
(subscription or payment may be required)

Copyright © 2019 [HealthDay](#). All rights reserved.

(HealthDay)—For postmenopausal women, the prevalence of knee osteoarthritis is lower with menopausal hormone therapy (MHT) use, according to a study published online Dec. 21 in *Menopause*.

Jae Hyun Jung, M.D., Ph.D., from the Korea University College of Medicine in Seoul, and colleagues collected data from 4,766 [postmenopausal women](#) from the Korea National Health and Nutrition Examination Survey (2009 to 2012). MHT was defined as use of regular hormone medication for one year or more; the correlation between prevalence of [knee](#) OA and MHT was examined.

The researchers found that the odds ratio for OA was 0.7 for the MHT group versus the non-MHT group in the multiple logistic regression models.

"This cross-sectional study precludes conclusions about causal relationships, so further prospective studies and intervention trials should be

APA citation: Prevalence of knee OA lower with menopausal hormone therapy (2019, January 17) retrieved 19 October 2019 from <https://medicalxpress.com/news/2019-01-prevalence-knee-oa-menopausal-hormone.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.