

Siberian indigenous people tolerate tick-borne infections more easily

February 5 2019, by Elena Fritz



Nikolay Ilyinsky, Dr.Sci.Biol, scientist at the TSU Biological Institute and Siberian State Medical University. Credit: Tomsk State University

Tomsk scientists have conducted a study to identify the characteristics of

the course of tick-borne infections and their consequences among residents of several Siberian regions. The findings suggest that the indigenous peoples of the North, and the sixth- and eighth-generation descendants of Siberians more easily tolerate, and recover more quickly from, infection with tick-borne encephalitis (TBE), tick-borne borreliosis (Lyme disease), and human granulocytic anaplasmosis (HGA).

"Under the project, [patients](#) from Tomsk, Surgut, Tyumen, and Khanty-Mansiysk were examined. Colleagues from these cities helped us a lot with their selection," says Nikolay Ilyinsky, Dr.Sci.Biol, scientist at the TSU Biological Institute and Siberian State Medical University.

"We have shown that in the case of tick-borne infections in humans, genetically transformed [cells](#) appear. In the future it will be an unfavorable sign, so the task of our study was to determine the time periods at which people who have had tick-borne infections normalize the number of cells with genetic changes to the level characteristic of a healthy person. In this connection, we estimated the number of cytogenetic changes by analyzing the number of cells with micronuclei in the buccal epithelium (on the inside of the cheek), leukocytes, and erythrocytes of human blood."

It was found that the proportion of cells with [chromosomal abnormalities](#) in patients with TBE, Lyme disease, or HGA increases sharply at the height of the illness (the first week of hospitalization). In some patients, the full normalization of the level of cells with an abnormal structure is observed in a month. In 39.8% of cases, this occurs after 3 months; 12.6% of patients have it after 6 months, and in 5.2% of cases, it does not decrease to the desired level even after that.

Patients who have had tick-borne borreliosis often have a complication in the form of arthritis, most often affecting the knee joints. Borrelia

settle in the cartilage tissue and destroy it, which ultimately leads to the erosion of cartilage, bone deformation, and arthrosis. In the study of the articular fluid of patients with Lyme disease, scientists identified cells with an abnormal structure, including those with extra chromosome number 7 (in total, instead of 46 chromosomes, 47 chromosomes were present in the transformed cells of a sick person). It is noted that the presence of this pathology exacerbates the patient's degree of motor impairment.

"An interesting fact is that certain groups of patients who have had tick-borne [borreliosis](#) show [high resistance](#) to such adverse effects," says Nikolay Ilyinsky. These are the indigenous people of Siberia—Khanty and Mansi, and Russians who are Siberians in the sixth or eighth generation, that is, people who are well adapted to the conditions of Siberia. The reason for such stability is not known; maybe it is a kind of mechanism of natural selection. There is another point of view: our Novosibirsk colleagues suggest that the whole thing is in the innate immunity that the child acquires from the mother (especially in the rural population).

Perhaps for the same reason, representatives of the above groups are more likely to carry tick-borne infections. The severe course of the disease with chronicity of the process is noted in people who came from the southern regions of the European part of Russia and the CIS c

Along with this, scientists have established a number of factors contributing to the long-term preservation of elevated levels of cells with chromosomal changes: these are alimentary obesity in women and the presence of type II diabetes in men and women. Particularly high and long-lasting preservation of an increased number of cytogenetically transformed cells is observed in mixed infections of tick-borne encephalitis, Lyme disease, and HGA.

The project scientists were the first to obtain data on patients with HGA. Currently world literature provides information only on animals. In the Tomsk Region, this tick-borne infection transmitted to humans has been observed only in the last seven or eight years. The chromosomal analysis of peripheral blood lymphocytes in patients with HGA showed an increase in the number of cells with abnormalities in chromosome 18 and 21. Along with this, in men aged 24-45 years, there was an increased frequency of deviations on the same chromosomes in the spermatozoa, which can have a negative effect on the person's reproductive functions.

Professor Ilinsky notes that the results are intended primarily for medical and genetic services and practicing infectious disease doctors. In addition, the new data can be used by obstetrician-gynecologists in treating patients with tick-borne infections.

Provided by Tomsk State University

Citation: Siberian indigenous people tolerate tick-borne infections more easily (2019, February 5) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2019-02-siberian-indigenous-people-tolerate-tick-borne.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--