

What is mastitis?

14 February 2019, by Leanda Mckenna, Adelle Mcardle And Kathryn Shine



Mastitis can be extremely painful. Credit: shutterstock.com

It's the middle of the night. Your newborn baby is awake. Again. She wants to feed. You lift her to your breast and brace for the pain.

Most [mothers](#) who breastfeed come to appreciate the convenience and the bonding it provides. But until your breasts get used to feeding your infant, it may hurt.

Around [one in five Australian mothers](#) will also develop [mastitis](#), an inflammation of the breast tissue. Mastitis is most common in the first [four to six weeks](#) after birth.

How do you know you've got mastitis?

[Common symptoms](#) of mastitis include a breast that may be abnormally red, tender to touch, or painful. You may have swelling, firm areas or lumps in your breast.

Some mothers may also have more general flu-like symptoms such as a fever, chills, high temperature, fatigue, joint aches, headache or malaise.

Your [health care](#) professional will typically diagnose mastitis based on these symptoms; blood tests aren't usually needed.

Mastitis can be [very distressing](#) for [new mothers](#), as it impacts on their desire to continue breastfeeding, their capacity to cope with their baby, and in some cases, their ability to care for other children.

About 10% of breastfeeding mothers get [mastitis more than once](#) when breastfeeding the same baby, and 7% will get mastitis again when breastfeeding subsequent children.

The [more children a woman has](#), the less likely she is to develop mastitis during breastfeeding. This may be because women become more used to breastfeeding, the more children they have.

Causes

[Nipple damage](#) may lead to mastitis by allowing bacteria to enter the breast tissue through a graze or crack on the nipple. Cracked nipples can occur if the baby has difficulty attaching to the breast during breastfeeding, or when the baby's feeding action damages the skin of the nipple.

About 60% of mothers who have mastitis have an [infection](#) caused by the bacteria that commonly live on the skin or in the breast. However, these same bacteria are present in the milk of mothers who don't have mastitis. So although a [bacterial infection](#) may be the cause of mastitis for some, it's not necessarily the cause for all mothers with the condition.

Milk stasis, or the obstruction of milk flow, can also be an important factor in the development of mastitis. Breast milk needs to be [effectively cleared](#) from the breast and there are a number of reasons why this may not be the case.

Mothers may have an [oversupply of milk](#) or have begun to decrease their level of breastfeeding.

Wearing a bra with straps that cut across breast tissue may reduce milk flow, and clearance, from that section of the breast.

Using dummies to [soothe](#) infants who may be hungry can delay breastfeeding and subsequent milk clearance from the breast.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

If the breast is not cleared of milk, milk stasis may allow bacterial concentrations to change within it, or cause an inflammatory reaction as milk is forced out of milk ducts and into the surrounding breast tissue.

Treatment and prevention

Effectively removing milk from the breast is the [most important](#) part of treatment for mastitis. This can usually be achieved by helping the baby attach properly to the breast – with the assistance of a lactation consultant – and by feeding regularly.

Provided by The Conversation

To manage your symptoms, your health care professional might suggest taking pain relief, resting, drinking lots of water, applying warmth to help with milk clearance (such as by having a warm shower), and applying cooling after [breastfeeding](#).

GPs may prescribe antibiotics to treat mastitis, but there is [very little evidence](#) to show this approach is effective.

Physiotherapists can treat mastitis using [ultrasound and gentle massage](#) to help remove [milk](#) from the breast. [Research](#) is underway to evaluate how well used and effective these emerging techniques may be.

In 3% of mothers with mastitis, it may progress to a [breast abscess](#), which may require hospitalisation and [treatment](#) via needle aspiration. In serious cases, the condition may require surgical drainage.

There is now also some evidence to suggest that [probiotics such as Lactobacillus](#) may be an effective preventive measure. They have been shown to halve the chance of developing mastitis when taken for 16 weeks following childbirth.

To avoid mastitis, mothers wearing maternity bras should ensure the bra does not cut across the [breast](#) tissue. If you're using dummies, only give them when you're sure the baby isn't hungry. Most importantly, continue to breastfeed regularly.

APA citation: What is mastitis? (2019, February 14) retrieved 19 September 2019 from <https://medicalxpress.com/news/2019-02-mastitis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.