

# Scientists identify unique subtype of eczema linked to food allergy

20 February 2019



A researcher demonstrates the minimally invasive collection of skin samples using small, clear tape strips. Credit: National Jewish Health

Atopic dermatitis, a common inflammatory skin condition also known as allergic eczema, affects nearly 20 percent of children, 30 percent of whom also have food allergies. Scientists have now found that children with both atopic dermatitis and food allergy have structural and molecular differences in the top layers of healthy-looking skin near the eczema lesions, whereas children with atopic dermatitis alone do not. Defining these differences may help identify children at elevated risk for developing food allergies, according to research published online today in *Science Translational Medicine*. The research was supported by the National Institute of Allergy and Infectious Diseases (NIAID), part of the National Institutes of Health.

"Children and families affected by food allergies must constantly guard against an accidental exposure to foods that could cause life-threatening allergic reactions," said NIAID Director Anthony S. Fauci, M.D. "Eczema is a risk factor for developing

food allergies, and thus [early intervention](#) to protect the [skin](#) may be one key to preventing [food allergy](#)."

Children with [atopic dermatitis](#) develop patches of dry, itchy, scaly skin caused by allergic inflammation. Atopic dermatitis symptoms range from minor itchiness to extreme discomfort that can disrupt a child's sleep and can lead to recurrent infections in scratched, broken skin.

The study, led by Donald Y.M. Leung, M.D., Ph.D., of National Jewish Health in Denver, examined the top layers of the skin, known as the stratum corneum, in areas with eczema lesions and in adjacent normal-looking skin. The study enrolled 62 children aged 4 to 17 who either had atopic dermatitis and [peanut allergy](#), atopic dermatitis and no evidence of any food allergy, or neither condition. Investigators collected skin samples by applying and removing small, sterile strips of tape to the same area of skin. With each removal, a microscopic sublayer of the first layer of skin tissue was collected and preserved for analysis. This technique allowed researchers to determine the skin's composition of cells, proteins and fats, as well as its microbial communities, [gene expression](#) within skin cells and water loss through the skin barrier.




A drawing of skin and a hair follicle. In this study, scientists analyzed the outermost layer of the skin, the stratum corneum, highlighted here in yellow. Credit: NIAMS, NIH

Researchers found that the skin rash of children with both atopic dermatitis and food allergy was indistinguishable from the skin rash of children with atopic dermatitis alone. However, they found significant differences in the structure and molecular composition of the top layer of non-lesional, healthy-appearing skin between children with atopic dermatitis and food allergy compared with children with atopic dermatitis alone. Non-lesional skin from children with atopic dermatitis and food allergy was more prone to [water loss](#), had an abundance of the bacteria *Staphylococcus aureus*, and had gene expression typical of an immature skin barrier. These abnormalities also were seen in skin with active atopic dermatitis lesions, suggesting that skin abnormalities extend beyond the visible lesions in children with atopic dermatitis and food allergy but not in those with atopic dermatitis alone.

"Our team sought to understand how healthy-looking skin might be different in children who develop both atopic dermatitis and food allergy compared to children with atopic dermatitis alone," said Dr. Leung. "Interestingly, we found those differences not within the skin rash but in samples of seemingly unaffected skin inches away. These insights may help us not only better understand atopic dermatitis, but also identify children most at risk for developing food allergies before they develop overt skin rash and, eventually, fine tune prevention strategies so fewer children are affected."

Allergy experts consider atopic dermatitis to be an early step in the so-called "atopic march," a common clinical progression found in some [children](#) in which atopic dermatitis progresses to food allergies and, sometimes, to respiratory allergies and allergic asthma. Many immunologists hypothesize that food allergens may reach immune cells more easily through a dysfunctional skin barrier affected by [atopic](#) dermatitis, thereby setting off biological processes that result in food allergies.

**More information:** D.Y.M. Leung et al., "The nonlesional skin surface distinguishes atopic dermatitis with food allergy as a unique endotype," *Science Translational Medicine* (2019). [stm.sciencemag.org/lookup/doi/10.1126/scitranslmed.aav2685](http://stm.sciencemag.org/lookup/doi/10.1126/scitranslmed.aav2685)



Provided by NIH/National Institute of Allergy and Infectious Diseases

The integrity of the skin barrier (arrows) is abnormal in the skin of atopic dermatitis patients with food allergy (A) compared to the skin of patients without food allergy (B) and controls (C). Credit: D.Y.M. Leung et al., *Science Translational Medicine* (2019)

APA citation: Scientists identify unique subtype of eczema linked to food allergy (2019, February 20)  
retrieved 16 June 2019 from <https://medicalxpress.com/news/2019-02-scientists-unique-subtype-eczema-linked.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*