

Why you lose hearing for a while after listening to loud sounds

8 March 2019



Anders Fridberger, professor at Linköping University.
Credit: Thor Balkhed

Loud sounds can impair hearing for a short time. Researchers at Linköping University, Sweden, have discovered a mechanism that explains how this happens. Their results are presented in the journal *Proceedings of the National Academy of Sciences, PNAS*.

"Most people have experienced [hearing impairment](#) and ear numbness after loud sounds. After a while, [hearing](#) returns to normal. We have discovered that a tiny structure in the cochlea known as the tectorial [membrane](#) plays an important role in this process, by acting as a storage depot for [calcium ions](#). These [calcium](#) ions contribute to regulating the function of the sensory cells," says Anders Fridberger at the Department of Clinical and Experimental Medicine, who led the study.

Calcium ions, which are calcium atoms with a positive charge, play a key role in the processes that make hearing possible. The conversion of sound waves to nerve impulses takes place in the inner ear, also known as the cochlea, which looks

like a spiral snail shell. The cochlea contains many sensory cells that detect sounds and generate signals that are passed on to the brain.

Previous research has shown that the fluid surrounding the sensory cells in the cochlea has a low concentration of calcium ions. There were, however, questions surrounding this, because sensory cells that are placed in fluids with the natural level of calcium no longer work normally.

The scientists at LiU have investigated the calcium ion concentration in the inner ear of guinea pigs, which is very similar to the human ear. By fluorescently labeling the calcium ions, the researchers discovered that a membrane that lies on top of the sensory cells, the tectorial membrane, has a much higher concentration of calcium ions than the surrounding fluid. The membrane seems to function as a store, such that the sensory cells are surrounded by higher levels of calcium ions than previously believed.

When the scientists added a substance that mops up calcium ions, the sensory cells ceased to function. In the next step, they exposed the inner ear to noise levels that correspond to those experienced at rock concerts, which had the same effect.

"When we expose isolated inner ears to [loud noise](#) in the lab, the level of calcium in the tectorial membrane falls, and the sensory [cells](#) cease to function. After a while, however, the calcium ion concentration returns to its previous level, and the [sensory cells](#) start to function again," says Anders Fridberger.

Until now, scientists have thought that the tiny membrane had a mainly mechanical function, but the discovery made by the LiU researchers suggests that the tectorial membrane may play an important role in regulating hearing by storing calcium.

"We knew that the tectorial membrane is necessary for hearing, and that it must be intact and correctly located, but it has been unclear why damage to this membrane contributes to impaired hearing," says Pierre Hakizimana, one of the researchers behind the study.

The research group is now planning to investigate whether the same mechanism is important in age-related hearing impairment.

More information: Clark Elliott Strimbu et al. Control of hearing sensitivity by tectorial membrane calcium, *Proceedings of the National Academy of Sciences* (2019). DOI: [10.1073/pnas.1805223116](https://doi.org/10.1073/pnas.1805223116)

Provided by Linköping University

APA citation: Why you lose hearing for a while after listening to loud sounds (2019, March 8) retrieved 15 October 2019 from <https://medicalxpress.com/news/2019-03-loud.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.