


Stress hormones promote breast cancer metastasis

14 March 2019



Cancer cell during cell division. Credit: National Institutes of Health

It has long been thought that stress contributes to cancer progression. Scientists from the University of Basel and the University Hospital of Basel have deciphered the molecular mechanisms linking breast cancer metastasis with increased stress hormones. In addition, they found that synthetic derivatives of stress hormones, which are frequently used as anti-inflammatory in cancer therapy, decrease the efficacy of chemotherapy. These results come from patient-derived models of breast cancer in mice and may have implications for the treatment of patients with breast cancer, as the researchers report in the scientific journal *Nature*.

One major obstacle in the treatment of metastatic breast cancer is the phenomenon of tumor heterogeneity. As the disease progresses, the tumor becomes more diverse, and the difference between the cancer cells may lead to inadequate treatment.

Because the underlying mechanisms of this phenomenon remain unclear, the research group of Prof. Mohamed Bentires-Alj from the Department of Biomedicine at the University of Basel and University Hospital of Basel has been studying the cells of a highly metastatic form of cancer known as triple-negative breast cancer. This cancer type is resistant to standard therapies leaving patients with fewer treatment options.

Stress accelerates metastasis

To explore the heterogeneity between tumors and metastases, the researchers profiled the activity of genes in a mouse model of breast cancer. They found that metastases have increased activity of glucocorticoid receptors (GR) which mediate the effects of stress hormones such as cortisol.

Concentrations of the stress hormones cortisol and corticosterone were higher in mice with metastases than in those with no metastases. The scientists show that increased levels of these [stress hormones](#) activate the GR, which cause increased colonization and heterogeneity of the [cancer cells](#)—and ultimately, shortened survival.

Reduced efficacy of chemotherapy

GR also mediates the effects of synthetic derivatives of cortisol such as dexamethasone which is used widely to treat the side effects of chemotherapy. The research group shows that in mice with metastatic cells the efficacy of the chemotherapy drug paclitaxel was decreased when administered in combination with dexamethasone.

These findings suggest that caution should be taken when prescribing glucocorticoid hormones to patients with breast cancer. The study also suggests that GR inhibition may be beneficial for patients and could lead to the development of new therapies to combat [breast cancer metastasis](#).

"Tumor heterogeneity is a serious hurdle for therapy. These findings highlight the importance of stress management in patients—and especially those with [triple-negative breast cancer](#)," states Prof. Bentires-Alj. "Moderate exercise and relaxation techniques have been shown to correlate with enhanced quality of life and greater survival in patients."

More information: Glucocorticoids promote breast cancer metastasis, *Nature* (2019). DOI: [10.1038/s41586-019-1019-4](https://doi.org/10.1038/s41586-019-1019-4), <https://www.nature.com/articles/s41586-019-1019-4>

Provided by University of Basel

APA citation: Stress hormones promote breast cancer metastasis (2019, March 14) retrieved 20 March 2019 from <https://medicalxpress.com/news/2019-03-stress-hormones-breast-cancer-metastasis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.