

Shift work increases diabetes and heart disease risk

3 April 2019



Credit: CC0 Public Domain

With over 20% of the population in industrial countries engaging shift work—in sectors such as healthcare and transportation—we urgently need to understand its health burden.

Many studies have shown that shift work is associated with heart and <u>metabolic diseases</u>, but new research in *Experimental Physiology* has clarified how shift work can have a long-term effect on the risk of heart disease and diabetes.

The study specifically suggested that shift work has a <u>negative impact</u> on the way a type of fat (called triglycerides) is broken down, as well as on the way sugar is utilised in our bodies. Both of these increase the risk of heart disease and diabetes because they affect how our body processes sugar and fat.

Researchers at the University College of Medical Sciences, University of Delhi conducted the research on two groups of healthcare workers. The first group included nurses, doctors and other healthcare workers aged 20 to 40 of both sexes who had not done <u>night shift</u> in the last one year or ever and had normal blood sugar levels. The

second group was of the same professional background and age, but involved in rotational night shift duties (more than 4 nights duties per month at least for last one year) and had normal blood sugar levels.

Blood <u>sugar</u> levels were measured using an oral glucose tolerance test. Then, after 12 hours of overnight fasting, participants were given a high fat meal. Fasting insulin levels, and triglyceride levels, after fasting and after the meal, were measured in all of the <u>study participants</u>. These were compared between health care workers with and without night shift duties.

Lead author on the study, SV Madhu said:

"This study gives us a better understanding of why shift work is associated, in the long-term, with heart and metabolic diseases, helping us work towards reducing the incidence of heart disease, diabetes and obesity in the future."

More information: Keithellakpam Kiranmala et al, Association of postprandial triglyceride responses with insulin resistance among rotational night shift healthcare workers, *Experimental Physiology* (2019). DOI: 10.1113/EP087514

Provided by The Physiological Society



APA citation: Shift work increases diabetes and heart disease risk (2019, April 3) retrieved 18 September 2019 from <u>https://medicalxpress.com/news/2019-04-shift-diabetes-heart-disease.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.