

New formula better predicts speed of tumor growth in 12 cancers

April 3 2019, by Marcene Robinson

University at Buffalo researchers have developed a new method to more accurately predict tumor growth rates, a crucial statistic used to schedule screenings and set dosing regimens in cancer treatment.

The [mathematical method](#) successfully estimated the doubling time—the amount of time for a tumor to double in size—for 12 types of cancer, ranging from breast and [prostate cancers](#) to melanoma.

The research, published in February in the *AAPS Journal*, was led by Dhaval Shah, Ph.D., associate professor in the UB School of Pharmacy and Pharmaceutical Sciences.

"This novel method allows clinicians and drug development scientists to use routinely-generated [clinical data](#) to infer doubling times of solid tumors. This parameter can be used to design individualized dosing regimens and develop reliable models for anticancer therapeutics," says Shah.

Tumor doubling time can significantly affect the outcome of anticancer therapy, but the rate is challenging to determine. Current methods calculate doubling time by measuring the size of a tumor at two points in time and assuming the cancer will grow at an exponential rate.

However, most doubling times calculated using this method are overestimated, and tiny changes in tumor size can make determining growth rates difficult.

The error impacts the ability of clinicians to schedule optimal follow-up screenings, set effective dosing regimens, and determine whether surgery, chemotherapy or radiation therapy is the best form of treatment.

The UB researchers instead base their method on data extracted from progression-free survival plots—the length of time during and after treatment that a cancer does not grow or spread.

Progression-free survival plots, explains Shah, inherently contain information that could help identify tumor growth rates.

The investigators examined data from 47 clinical trials that reported plots for any of 12 cancer types: melanoma; pancreatic, lung, prostate, gastric, colorectal and three forms of breast cancer; hepatocellular (liver) and renal cell (kidney) carcinoma; and glioblastoma multiforme (brain).

The cancer [growth rates](#) predicted by the researchers using progression-free survival plots were within close range to the reported actual tumor doubling times.

More information: Katherine Kay et al. Estimation of Solid Tumor Doubling Times from Progression-Free Survival Plots Using a Novel Statistical Approach, *The AAPS Journal* (2019). [DOI: 10.1208/s12248-019-0302-5](#)

Provided by University at Buffalo

Citation: New formula better predicts speed of tumor growth in 12 cancers (2019, April 3) retrieved 23 April 2024 from <https://medicalxpress.com/news/2019-04-formula-tumor-growth-cancers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.