


# Researchers find tissue environment plays important role in tumor formation

April 26 2019, by Kathleen Haughney

---



Credit: CC0 Public Domain

A team of Florida State University researchers used a simple tumor model to show the importance of tissue microenvironment in primary tumor formation.

In the journal *Developmental Cell*, FSU researchers explain how certain signaling pathways define [tumor](#) hotspots, tissue microenvironment favorable for tumorigenesis, and show that simple stimulation of a protein can lead directly to tumor formation.

"The beauty of this tumor model is its simplicity and reproducibility," said Professor of Biological Science Wu-Min Deng. "We hope to use this model to understand how tumors initiate, progress and metastasize, and use it for drug screens as well."

Using a fruit fly model, Deng and former FSU graduate student Sheng-An Yang examined transition zones. Transition zones are areas where one type of epithelial cell meets another type and are often prone to tumors. For this study, researchers looked at a [transition zone](#) within the salivary gland imaginal ring—the area where two types of larval [salivary gland cells](#) meet.

The researchers found that when a signaling molecule called Notch is genetically stimulated in this transition zone—something that could also occur naturally by exposure to mutagens such as chemicals and radiation—it consistently turned a healthy cell into a cancerous one.

In fruit fly imaginal rings, these Notch proteins are required for cell proliferation. But in a mammal, Notch proteins perform numerous functions, including neural developmental and intestinal stem cell

maintenance.

"Cancer development has long been viewed as a complex multistep process that requires accumulation of oncogenic and tumor suppressor mutations," Yang said. "Our findings reveal an alternative route, where in tumor hotspots such as transition zones, overexpression of a type of gene that can potentially turn a cell into a tumor is sufficient to induce malignant tumor transformation."

Deng and Yang performed all of their research on [fruit flies](#). Fruit flies are used as a model to determine the basic fundamentals of many diseases in humans, including [cancer](#), because about 75 percent of human disease-causing genes have a fruit fly counterpart.

"Our experiments show that the imaginal ring tumors induced by upregulation of Notch eventually become malignant," Yang said.

Yang noted that their model doesn't necessarily cover all types of cancer, but it is a good starting point to look at other transition zones and specific kinds of cancerous tumor formations.

"Cancer is so complex and highly dynamic. It is hard to use a single model to cover all cancer types," he said. "Nevertheless, our study is still related to a broad range of cancer types, especially site-specific cancer formation."

Provided by Florida State University

Citation: Researchers find tissue environment plays important role in tumor formation (2019, April 26) retrieved 18 April 2024 from <https://medicalxpress.com/news/2019-04-tissue-environment-important-role-tumor.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.