

First in-vivo trial of subharmonic contrast-enhanced imaging for detection of PCa

6 May 2019

A new technique for imaging of microbubble ultrasound contrast agents may be useful in detection of prostate cancer (PCa) not found by multiparametric magnetic resonance imaging (MRI), according to a study to be presented at the ARRS 2019 Annual Meeting, set for May 5-10 in Honolulu, HI.

The first in vivo application of contrast-enhanced SHI in the prostate, the [pilot study](#) was conducted to evaluate contrast-enhanced subharmonic imaging (SHI) of the prostate for detection of PCa.

Building on the authors' previous work demonstrating the effectiveness of contrast enhanced harmonic imaging (HI) for detection of prostate cancer, 55 patients referred for [prostate biopsy](#) were imaged using conventional grayscale, color, and power Doppler, conventional contrast HI, SHI—a [new technique](#) for imaging of microbubble ultrasound contrast agents with up to a 10-fold increase in contrast-to-background signal ratio relative to conventional HI—and flash replenishment in combination with SHI (MIP-SHI).

The results demonstrated contrast-enhanced SHI enhancement in all patients. Detection of PCa using contrast-enhanced SHI included 9 of 31 patients with a prior negative MRI or negative MRI-guided biopsy, suggesting SHI may be useful in detection of PCa not found by multiparametric MRI.

"Diagnosis of clinically significant prostate cancer with non-invasive means is a real clinical challenge," said author of the study Ethan Halpern, MD. " Although multi-parametric MRI is currently used for this purpose, ultrasound has numerous advantages. Ultrasound systems are portable. Ultrasound studies are far less expensive and more widely available as compared with MRI. Contrast enhanced ultrasound studies can be performed at the same sitting as the ultrasound-guided biopsy, while mp-MRI requires two visits, one for the diagnostic MR study and a second for

the fusion biopsy. There is no need for a fusion imaging system (additional hardware/software) when using contrast enhanced ultrasound. No additional effort is required to properly register the contrast-enhanced ultrasound diagnostic study with the targeting system for biopsy. Combining contrast enhanced [ultrasound](#) with subharmonic imaging has the potential to provide a new, non-invasive method for diagnosis and characterization of [prostate](#) cancer."

More information: www.arrs.org/am19

Provided by American Roentgen Ray Society

APA citation: First in-vivo trial of subharmonic contrast-enhanced imaging for detection of PCa (2019, May 6) retrieved 19 November 2019 from <https://medicalxpress.com/news/2019-05-in-vivo-trial-subharmonic-contrast-enhanced-imaging.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.