

Significant unmet mental health care needs exist in current and former smokers with COPD

8 May 2019



Credit: CC0 Public Domain

Researchers at the University of Alabama at Birmingham have published new information that stresses the need for increased mental health care for current and former smokers, especially those who suffer from chronic obstructive pulmonary disease.

Published in the *Journal of Psychosomatic Research*, physician scientists from the UAB Division of Pulmonary, Allergy and Critical Care Medicine in partnership with researchers across the country say [smokers](#) with and without COPD have significant unmet [mental health care](#) needs, particularly anxiety and depressive symptoms.

"In the group of more than 5,000 smokers with and without COPD from the multisite, longitudinal genetic epidemiology of COPD—COPDGene—study, one in four had clinically elevated anxiety/depressive symptoms," said study lead author and UAB assistant professor Anand S. Iyer, M.D. "Of those with elevated symptoms, two-thirds were not receiving

any type of medication to treat these symptoms such as an anti-anxiety or anti-depressant medication. There are many non-pharmacological treatment options, too; but medications are cornerstone therapy for clinically elevated symptoms."

Iyer added that these symptoms were not only a problem for smokers with end-stage COPD.

"While [depressive symptoms](#) were most frequent in those with the most severe cases of COPD, anxiety symptoms were similar in frequency between smokers with and without COPD, so the problem seems to start a lot earlier in the disease," Iyer said. "We also identified several characteristics that were associated with having unmedicated symptoms—namely African-American race, male gender and lack of health insurance."

According to Iyer, who is also part of the UAB Lung Health Center, untreated emotional symptoms leave a major gap in COPD care that warrants early [palliative care](#) or primary palliative care. He says emotional symptoms have been shown to impact many COPD-related outcomes.

"We have previously found that depression predicts hospital readmissions in COPD, and others have found associations with exacerbations and poor adherence to medications," Iyer said. "Clinically, we treat many COPD patients and smokers who have borderline COPD and also have clinically important emotional symptoms that would warrant treatment."

Iyer added that pulmonary and primary care clinicians do not assess these symptoms often enough, which leaves many patients potentially not being treated for elevated emotional symptoms, a major gap in comprehensive COPD care, and it is not clear who is in charge of managing those

symptoms—primary care or pulmonologists.

"Since the elevated emotional symptoms we discovered were also present in smokers who had milder COPD, earlier assessment and triage in the primary care setting is needed, potentially even early palliative care," he said.

So, does smoking cause anxiety or depression? Iyer says that relationship can be tricky, but seems to be bidirectional.

"There are similar receptors in the brain functioning in smokers addicted to cigarettes and who have anxiety and depression," Iyer said. "In fact, most of the smoking cessation medications frequently used to help patients quit are antidepressants too."

"Imagine someone who is anxious and self-medicating that anxiety with cigarettes, which we obviously don't want," he said. "There are many phenotypes in COPD, such as someone with chronic cough or emphysema or high eosinophil count, and all of these warrant unique patient-centered approaches. Our research adds to the growing literature on a possible unique COPD-psychiatric phenotype who is at high risk for exacerbations and poor quality of life."

Iyer says the next steps for this research are to see how symptoms of [anxiety](#) and depression influence important COPD outcomes, particularly exacerbations of COPD.

More information: Anand S. Iyer et al, Symptoms of anxiety and depression and use of anxiolytic-hypnotics and antidepressants in current and former smokers with and without COPD - A cross sectional analysis of the COPDGene cohort, *Journal of Psychosomatic Research* (2019). DOI: [10.1016/j.jpsychores.2019.01.002](https://doi.org/10.1016/j.jpsychores.2019.01.002)

Provided by University of Alabama at Birmingham
APA citation: Significant unmet mental health care needs exist in current and former smokers with COPD (2019, May 8) retrieved 21 September 2019 from <https://medicalxpress.com/news/2019-05-significant-unmet-mental-health-current.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.